

Cryptococcus gattii, Strain Alg75

Catalog No. NR-50187

Product Description: *Cryptococcus gattii* (*C. gattii*), strain Alg75 is the progeny of a genotypic cross between *C. gattii* strains R265 and Alg40.

Lot¹: 63910608

Manufacturing Date: 21DEC2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology ²	Report results	Globose to ovoid, single or budding (Figure 1A)
Colony morphology ³	Report results	Smooth, entire and white to cream (Figure 1B)
Canavanine-glycine-bromthymol blue (CGB) differential medium ⁴	Report results	Blue
Genotypic Analysis		
Sequencing of partial 18S rRNA gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 28S rRNA (~ 510 base pairs)	≥99% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1)	100% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1)
Sequencing of 26S rRNA gene (~ 620 base pairs)	≥99% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)	100% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)
Purity⁵		
Nutrient broth with 0.1% Yeast Extract at 25°C	No bacterial growth	No bacterial growth
Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth	No bacterial growth
Viability (post-freeze)²	Growth	Growth

¹NR-50187, lot 63910608 was produced by incubation of seed material in modified Sabouraud Dextrose medium and incubated for 3 days at 25°C in an aerobic atmosphere. Yeast were harvested from agar plates with 20% glycerol prior to vialing.


²4 days at 25°C in an aerobic atmosphere on modified Sabouraud Dextrose agar

³3 days at 25°C in an aerobic atmosphere on modified Sabouraud Dextrose agar

⁴4 days at 26°C in an aerobic atmosphere. CGB medium differentiates *C. gattii* from *C. neoformans* based on the ability of *C. gattii* isolates to grow in the presence of L-canavanine and to assimilate glycine as a sole carbon source, resulting in a blue color. *C. neoformans* isolates will show yellow to light-green on CGB medium. [McTaggart, L., et al. "Rapid Identification of *Cryptococcus neoformans* var. *grubii*, *C. neoformans* var. *neoformans*, and *C. gattii* by Use of Rapid Biochemical Tests, Differential Media, and DNA Sequencing." *J. Clin. Microbiol.* 2011 (49): 2522-2527. PubMed: 21593254.]

⁵Clarity of broth was determined by visual inspection after 7 days at 25°C and 37°C in an aerobic atmosphere.

Figure 1: Cellular and Colony Morphology



Certificate of Analysis for NR-50187

Date: 18 JUL 2016

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

