

Cryptococcus neoformans*, Strain KN99α*Catalog No. NR-48769****Product Description:** *Cryptococcus neoformans* (*C. neoformans*), strain KN99α was derived from strain H99O, after passage in the rabbit model of central nervous system infection.**Lot¹: 63383703****Manufacturing Date: 07APR2015**


TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology ²	Report results	Circular yeast form cells, usually single (Figure 1A)
Colony morphology ²	Report results	Circular, convex, butyrous, shiny, smooth; entire margin (Figure 1B)
CGB agar characterization ³		
NR-48769	Yellow (no color change)	Yellow (no color change)
Positive control (<i>C. neoformans</i> ; ATCC® 32045™)	Yellow (no color change)	Yellow (no color change)
Negative control (<i>C. gattii</i> ; ATCC® MYA-4563™)	Blue	Blue
Genotypic Analysis		
Sequencing of partial 18S ribosomal RNA (rRNA) gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 26S rRNA (~ 910 base pairs)	≥ 99% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)	99.9% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)
Sequencing of 26S rRNA gene (~ 600 base pairs)	≥ 99% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)	99.8% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)
Purity⁴		
Nutrient broth with 0.1% Yeast Extract at 25°C	No bacterial growth	No bacterial growth
Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth	No bacterial growth
Viability (post-freeze)²	Growth	Growth

¹NR-48769 was produced by inoculation of the deposited material onto Yeast Mold slants and grown 4 days at 30°C in an aerobic atmosphere. Cells were harvested from the slants with 20% glycerol to produce this lot.

²3 days at 25°C in an aerobic atmosphere on Modified Sabouraud Dextrose medium

³2 days at 35°C in an aerobic atmosphere. CGB medium differentiates *C. gattii* from *C. neoformans* based on the ability of *C. gattii* isolates to grow in the presence of L-canavanine and to assimilate glycine as a sole carbon source, resulting in a blue color. *C. neoformans* isolates will remain yellow. [McTaggart, L., et al. "Rapid Identification of *Cryptococcus neoformans* var. *grubii*, *C. neoformans* var. *neoformans*, and *C. gattii* by Use of Rapid Biochemical Tests, Differential Media, and DNA Sequencing." *J. Clin. Microbiol.* 2011 (49): 2522-2527. PubMed: 21593254.]

⁴Clarity of broth was determined by visual inspection after 6 days in an aerobic atmosphere.

Figure 1: Cellular and Colony Morphology

Date: 27 SEP 2016**Signature:**

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

