

***Cryptococcus neoformans*, Strain H99W**

**Catalog No. NR-48773**

**Product Description:** *Cryptococcus neoformans* (*C. neoformans*), strain H99W was derived from strain H99O, after an unknown number of laboratory passages.

**Lot<sup>1</sup>: 63383707**

**Manufacturing Date: 30MAR2015**

TEST	SPECIFICATIONS	RESULTS
<b>Phenotypic Analysis</b> Cellular morphology <sup>2</sup>  Colony morphology <sup>2</sup>  CGB agar characterization <sup>3</sup> NR-48773 Positive control ( <i>C. neoformans</i> ; ATCC® 32045™) Negative control ( <i>C. gattii</i> ; ATCC® MYA-4563™)	Report results  Report results  Yellow (no color change) Yellow (no color change) Blue	Circular yeast form cells, usually single (Figure 1A) Circular, convex, butyrous, shiny, smooth; entire margin (Figure 1B)  Yellow (no color change) Yellow (no color change) Blue
<b>Genotypic Analysis</b> Sequencing of partial 18S ribosomal RNA (rRNA) gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 26S rRNA (~ 910 base pairs) Sequencing of 26S rRNA gene (~ 600 base pairs)	≥ 99% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)  ≥ 99% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)	99.9% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)  99.8% sequence identity to <i>C. neoformans</i> , strain H99 (GenBank: CP003821)
<b>Purity<sup>4</sup></b> Nutrient broth with 0.1% Yeast Extract at 25°C Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth No bacterial growth	No bacterial growth No bacterial growth
<b>Viability (post-freeze)<sup>2</sup></b>	Growth	Growth


<sup>1</sup>NR-48773 was produced by inoculation of the deposited material onto Yeast Mold slants and grown 3 days at 30°C in an aerobic atmosphere. Cells were harvested from the slants with 20% glycerol to produce this lot.

<sup>2</sup>3 days at 25°C in an aerobic atmosphere on Modified Sabouraud Dextrose medium

<sup>3</sup>2 days at 35°C in an aerobic atmosphere. CGB medium differentiates *C. gattii* from *C. neoformans* based on the ability of *C. gattii* isolates to grow in the presence of L-canavanine and to assimilate glycine as a sole carbon source, resulting in a blue color. *C. neoformans* isolates will remain yellow. [McTaggart, L., et al. "Rapid Identification of *Cryptococcus neoformans* var. *grubii*, *C. neoformans* var. *neoformans*, and *C. gattii* by Use of Rapid Biochemical Tests, Differential Media, and DNA Sequencing." *J. Clin. Microbiol.* 2011 (49): 2522-2527. PubMed: 21593254.]

<sup>4</sup>Clarity of broth was determined by visual inspection after 6 days in an aerobic atmosphere.

**Figure 1: Cellular and Colony Morphology**



Date: 27 SEP 2016

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.  
You are authorized to use this product for research use only. It is not intended for human use.

