

Zika Virus, H/PAN/2015/CDC-259359

Catalog No. NR-50219

Product Description: Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells¹ infected with Zika virus (ZIKV), H/PAN/2015/CDC-259359

Passage History: V1/V3 (Centers for Disease Control/BEI Resources); V# = Number of passages in Vero cells

Lot²: 64158291

Manufacturing Date: 23MAR2016

TEST	SPECIFICATIONS	RESULTS
Infectivity in Vero E6 Cells¹	Report results	Cell rounding and detachment
Sequencing of Species-Specific Region (976 nucleotides)	Consistent with ZIKV	Consistent with ZIKV ³
Titer by TCID₅₀ Assay in Vero E6 Cells^{1,4,5}	Report results	1.6 × 10 ⁶ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

¹Vero 76, clone E6: ATCC® CRL-1586™

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate (ATCC® 30-2003) and 2% fetal bovine serum (ATCC® 30-2020) for 5 days at 37°C with 5% CO₂

³Partial nucleotide sequencing of NR-50219 obtained at the time this lot was manufactured is consistent with contemporary ZIKV isolates from the Americas. The complete genomic sequence of NR-50219, Lot No. 64158291 has since been determined and deposited at NCBI (GenBank: KX156774, ZIKV/Homo sapiens/PAN/CDC-259359_V1-V3/2015).

⁴The Tissue Culture Infectious Dose 50% (TCID₅₀) is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁵8 days at 37°C with 5% CO₂

⁶Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 15 MAR 2017

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

