

Sua E1, *Anopheles gambiae* Cell Line

Catalog No. MRA-924

For research use only. Not for human use.

Contributor:

George K. Christophides, Ph.D., Professor of Infectious Diseases and Immunity, Department of Life Sciences, Faculty of Natural Sciences, Imperial College London, London, United Kingdom

Manufacturer:

BEI Resources

Product Description:

The *Anopheles gambiae* cell line Sua E1 was established from neonate larvae of the Suakoko 2La strain.¹

Material Provided:

Each vial of MRA-924 contains approximately 0.5 mL of Sua E1 cells in Schneider's Insect medium supplemented with 10% fetal bovine serum (FBS) and 10% dimethylsulfoxide (DMSO). Please see Appendix I for media preparation. Sufficient cells are provided to initiate at least one new culture. The cell count, expressed as cells per vial, is shown on individual certificates of analysis for each product lot.

Packaging/Storage:

This product was packaged aseptically, in screw-capped plastic cryovials. It should be stored at cryogenic temperature (-100°C or colder), preferably in the vapor phase of a liquid nitrogen freezer. Storage at -70°C will result in loss of viability. To insure the highest level of viability, the vial should be thawed and the culture initiated as soon as possible upon receipt. Any warming of the product during shipping and transfer must be avoided, as this will adversely affect the viability of the product after thawing. For transfer between freezers and shipping, the cells may be placed on dry ice for brief periods, although use of a portable liquid nitrogen carrier is preferred. Please read the following recommendations prior to reconstituting this material.

Safety Precautions:

When handling frozen vials it is highly recommended that protective gloves, lab coat and full face mask be worn. Even brief exposure to the ultra-cold temperature can cause tissue damage from frostbite. Also, some vials may slowly fill with liquid nitrogen if they have been immersed during cryogenic storage. When thawing, the liquid nitrogen may rapidly expand as it changes to gas, breaking the vial or cap with explosive force, sending debris flying with enough velocity to cause injury. Store and use in areas with adequate ventilation.

Growth Conditions:

Prior to thawing the cells, prepare culture medium according to Appendix I. Thaw 1 vial in a 25°C water bath and transfer

the contents into a 25-mL vented cell culture flask with 9 mL of culture medium. Keep the flask tightly capped in a 25°C incubator (no CO₂ required). Change the media at 12 to 16 hours post seeding. Feed the cells at least every 48 hours, harvest at 80%-90% confluency and reseed at a 1:3 to 1:5 ratio.

Sub-culture procedure: When cells near confluency, detach cells by vigorous shaking, mechanical disruption or gentle cell scraping. Collect and gently aspirate several times with a pipette to disrupt clumped cells prior to cell counting as required and passage to new flasks.

Note: Trypsin or trypsin-like enzyme substitute may be used to fully disperse adherent cells but is not recommended on a continuous basis for MRA-924.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Sua E1, *Anopheles gambiae* Cell Line, MRA-924, contributed by George K. Christophides."

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370

Fax: 703-365-2898

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Christophides, G. K., Personal Communication.
2. Müller, H.-M., et al. "A Hemocyte-Like Cell Line Established from the Malaria Vector *Anopheles gambiae* Expresses Six Prophenoloxidase Genes." J. Biol. Chem. 274 (1999): 11727-11735. PubMed: 10206988.
3. Dimopoulos, G., et al. "Genome Expression Analysis of *Anopheles gambiae*: Responses to Injury, Bacterial Challenge, and Malaria Infection." Proc. Natl. Acad. Sci. USA 99 (2002): 8814-8819. PubMed: 12077297.

ATCC® is a trademark of the American Type Culture Collection.



APPENDIX I: MEDIA PREPARATION

Culture Medium

Schneider's Insect medium
10% FBS (qualified for insect cell culture or heat-inactivated)

Facultative:
100 U/mL Penicillin
100 U/mL Streptomycin

Freezing Medium

Schneider's Insect medium
10% FBS
10% DMSO