

***Plasmodium falciparum*, Strain IPC 5202**

Catalog No. MRA-1240

Product Description: *Plasmodium falciparum* (*P. falciparum*), strain IPC 5202 was isolated in 2011 from a human patient with malaria in Battambang province, western Cambodia. *P. falciparum*, strain IPC 5202 has shown resistance to artemisinin.

Lot¹: 62401478

Manufacturing Date: 24FEB2014

TEST	SPECIFICATIONS	RESULTS
Identification by Giemsa Stain Microscopy²	Blood-stage parasites present	Blood-stage parasites present
Antimalarial Susceptibility Profile (<i>in vitro</i>) Half-maximal Inhibitory Concentration (IC ₅₀) by SYBR green I [®] drug sensitivity assay ³ Chloroquine Artemisinin Quinine Cycloguanil Pyrimethamine Sulfadoxine Ring-stage Survival Assay (RSA _{0-3h}) ⁴ Dihydroartemisinin (DHA) ⁵	Report results Report results Report results Report results Report results Report results Report results	18.8 ± 2.2 nM 26.9 ± 6.2 nM 165.5 ± 26.8 nM 670.4 ± 46.3 nM 30500 ± 4937.3 nM 532000 ± 61384.2 nM
Genotypic Analysis Sequencing of Merozoite Surface Protein 2 (MSP2) gene (~ 850 base pairs) MSP2 PCR amplicon analysis ⁶	Consistent with <i>P. falciparum</i> ~ 600-900 base pair amplicon	Consistent with <i>P. falciparum</i> (Figure 1) ~ 900 base pair amplicon
Level of Parasitemia Pre-freeze ⁷ Ring-stage parasitemia Total parasitemia Post-freeze ⁸ Ring-stage parasitemia Total parasitemia	Report results ≥ 2% Report results ≥ 1%	3.19% 4.78% 0.86% 1.50%
Viability (post-freeze)⁹	Growth in infected red blood cells	Growth in infected red blood cells
Sterility (21-day incubation) Harpo's HTYE broth ¹⁰ , 37°C and 26°C, aerobic Tryptic Soy broth, 37°C and 26°C, aerobic Sabouraud Dextrose broth, 37°C and 26°C, aerobic DMEM with 10% FBS, 37°C, aerobic Sheep Blood agar, 37°C, aerobic Sheep Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination DNA Detection by PCR	None detected	None detected

¹MRA-1240 was produced by cultivation of the deposited material in fresh human erythrocytes suspended in RPMI 1640 medium, adjusted to contain 10% (v/v) heat-inactivated human serum (pooled Type A), 25 mM HEPES, 2 mM L-glutamine, 4 g/L D-glucose, 0.005 µg/mL hypoxanthine and 2.5 µg/mL gentamicin. The culture was incubated at 37°C in sealed flasks outgassed with blood-gas atmosphere (90% N₂, 5% CO₂, 5% O₂) and monitored for parasitemia daily for 26 days. Every 1 to 4 days, uninfected, leukocyte filtered, Type O erythrocytes in complete culture medium were added dropwise to the culture as needed and monitored for hematocrit.

²Blood-stage malaria parasites (rings, trophozoites, schizonts +/- gametocytes) were examined by microscopic Giemsa-stained blood smears of an *in vitro* human blood culture over 3 days.

³A SYBR Green I[®] anti-malarial drug sensitivity assay in 96-well plates was used to determine IC₅₀ values of an active (> 70% ring stage) parasite culture in the presence of each antimalarial drug [Hartwig, C. L., et al. "XI: I. SYBR Green I[®]-Based Parasite Growth Inhibition Assay for Measurement of Antimalarial Drug Susceptibility in *Plasmodium falciparum*." In *Methods in Malaria Research Sixth Edition*. (2013) Moll, K., et al. (Ed.), EVIMalaR, pp. 122-129. Available at: <https://www.beiresources.org/Publications/MethodsInMalariaResearch.aspx>].

⁴A detailed RSA_{0-3h} protocol is available on the Worldwide Antimalarial Resistance Network's website at <http://www.wwarn.org/tools-resources/procedures/ring-stage-survival-assays-rsa-evaluate-vitro-and-ex-vivo-susceptibility>.

⁵*P. falciparum*, strain IPC 5202 was deposited in 2013 with a DHA RSA_{0-3h} value of 88.2%.

⁶Primer sequences and conditions for PCR are available upon request.

⁷Pre-freeze parasitemia was determined after 26 days post infection by microscopic counts of Giemsa-stained blood smears.

⁸Post-freeze parasitemia was determined after 3 days post infection by microscopic counts of Giemsa-stained blood smears.

⁹Viability was confirmed by examination of infected erythrocytes for parasitemia at 3 days post infection.

¹⁰Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Figure 1: MRA-1240 MSP2 Sequence

```
AACAACAAAT TTATTTATTG AAGCAATATT ACTAGAGTTA CTTAAGAGGG ATGTTGCTGC TCCACAGTTT TCTTTGTTAC
CATCGGTACA TTCTTTTTGA CTATCAGAAG TATTTTGTTG ATGATTATTT CTAGAACCAT GCATATGTCC ATGTTGTCCT
GTACCTTTAT TCTCTGGTGC AGATTGTAAT TCGGGGGATT CAGTTTGTTT GGCTGTTGGA GCAGAATTTT CAGCTTGTTT
AGGTTGTGCA GTAGGACTTT TAGTGTCTGC ATCTTGAGTG GGTGGAACAT TTGATTTAGT TTGAGAGTCT TGTTGAACAT
TTGAGTTATT TTGAGTTTCT TTATTTGCTT GATTTGGTTT TTGAACTTCT CCTTTACCTT TTGGATTTGT TTCGGCATT
TTATGATTTG GATTTTCTGA AGAGGTAAGT GTAGATGCTT CTGCATCATT AGTAGTTGTG GTAGTTGTGG TAGTAGCGGG
AGTACTTGGA CTTCTCTCAG CATCTGCACC AGGATTAGCA CCATTACCAT CACCAGAACC AGCACTACCA CTACCACCAG
CACTACCACT ACCACCAGCA CTACCACTAC CACCAGCACT ACCACTACCA CCAGCACTAC CACTACCACC AGCACTACCA
CTACCACCAG CACTACCACT ACCACCAGCA CTACCACTAC CACCAGCACT ACCACTACCA CCAGCACTAC CACTACCACC
AGCACCAGTA GAAGGATTAC TTTCTTCCAT ACTTCTCCTT ATACTCATAT TATAAGCATT GTTTATGAAT GTGTTGCTAT
ATTTACTTTC ATTTTAAATA TTAAAGGTAA CAAAAATAAA GAAATTTATA ATA
```

Date: 30 APR 2017

Signature: 

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

