

Dengue Virus Type 3, UIS 117

Catalog No. NR-49708

Product Description: Cell lysate and supernatant from *Aedes albopictus* mosquito larval clone C6/36 cells¹ infected with dengue virus type 3 (DEN-3), UIS 117

Passage History: V1C1/C3 (Prior to deposit at BEI Resources/BEI Resources); V# = Number of passages in Vero cells; C# = Number of passages in C6/36 cells²

Lot³: 70003599

Manufacturing Date: 20JAN2017

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using C6/36 Cells¹	Report results	Cell rounding and sloughing
Identification by Indirect Fluorescent Antibody (IFA) Assay⁴	Fluorescence observed	Fluorescence observed
Sequencing of Species-Specific Region (873 nucleotides)	Consistent with DEN-3	Consistent with DEN-3 ⁵
Titer by TCID₅₀ Assay^{6,7} in C6/36 Cells¹ with IFA Readout⁸	Report results	1.6 × 10 ⁷ TCID ₅₀ per mL
Amplification of Dengue Virus Sequence by RT-PCR	~ 1000 bp amplicon	~ 1000 bp amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁹ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹*Aedes albopictus* clone C6/36 cells (ATCC® CRL-1660™)

²The first viral passage at BEI Resources was performed by polyethylenimine (Polyplus-transfection® SA jetPEI® 101-10)-mediated transfection of extracted viral nucleic acid in order to remove contaminating mycoplasma.

³Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 7 days at 28°C with 5% CO₂.

⁴Using Anti-Dengue Virus Complex Antibody (Millipore MAB 8705)

⁵Sequence information for DEN-3, UIS 117 is not available in the NCBI database; nucleotide sequence obtained for NR-49708, Lot No. 70003599 is ~ 99% identical to numerous contemporaneous South American DEN-3 strains.

⁶The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.


⁷13 days at 28°C and 5% CO₂

⁸Using Anti-Dengue Virus Complex Antibody (Millipore MAB 8705)

⁹Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Certificate of Analysis for NR-49708

Date: 11 AUG 2017

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

