

***Pseudomonas aeruginosa*, Strain Stone No. 130**
Catalog No. NR-31040
Product Description: *Pseudomonas aeruginosa* (*P. aeruginosa*), strain Stone no. 130 was isolated in or before 1971 from a burn patient at Grady Memorial Hospital in Atlanta, Georgia, USA.

Lot¹: 70002167
Manufacturing Date: 03FEB2017

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ² Motility (wet mount) VITEK [®] MS (MALDI-TOF)	Gram-negative rods Report results Report results Consistent with <i>P. aeruginosa</i>	Gram-negative rods Circular, raised, undulate, smooth and cream (Figure 1) Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile VITEK [®] (AST-GN81 Card) ³ Ampicillin Amoxicillin/clavulanic Acid Piperacillin/tazobactam Cefazolin Cefoxitin Ceftazidime Ceftriaxone Cefepime Meropenem Amikacin Gentamicin Tobramycin Ciprofloxacin Levofloxacin Tetracycline Nitrofurantoin Trimethoprim/sulfamethoxazole Etest [®] antibiotic test strips ⁴ Gentamicin ⁵ Ofloxacin ⁵ Rifampicin ⁶ Streptomycin ⁷ Trimethoprim/sulfamethoxazole ⁸	Report results Report results Report results Report results Report results Report results Report results Report results Report results Report results Resistant Report results Report results Report results Report results Report results Resistant Resistant Report results Sensitive Resistant Resistant Resistant Resistant	Resistant (≥ 32 µg/mL) Resistant (≥ 32 µg/mL) Sensitive (≤ 4 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 64 µg/mL) Sensitive (≤ 1 µg/mL) Resistant (≥ 16 µg/mL) Sensitive (= 2 µg/mL) Sensitive (= 1 µg/mL) Sensitive (= 8 µg/mL) Resistant (≥ 16 µg/mL) Resistant (≥ 512 µg/mL) Resistant (≥ 320 µg/mL) Resistant (= 256 µg/mL) Sensitive (= 1.5 µg/mL) 32 µg/mL 1024 µg/mL 32 µg/mL ⁹
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1530 base pairs) Digital DNA-DNA hybridization (dDDH) ¹⁰	≥ 99% sequence identity to <i>P. aeruginosa</i> , strain Stone no. 130 (GenBank: AQFN01000045.1) ≥ 70% for species identification	100% sequence identity to <i>P. aeruginosa</i> , strain Stone no. 130 (GenBank: AQFN01000045.1) <i>P. aeruginosa</i> (91.4%) ¹¹
Purity (post-freeze)¹²	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)²	Growth	Growth


¹NR-31040 was produced by inoculation of the deposited material into Nutrient broth and grown for 1 day at 37°C in an aerobic atmosphere. Broth inoculum was added to Nutrient agar kolles, which were grown for 1 day at 37°C in an aerobic atmosphere to produce this lot.

²1 day at 37°C in an aerobic atmosphere on Nutrient agar

³Minimum Inhibitory Concentration (MIC); MIC interpretation was determined using VITEK[®] 2 software version 07.01 combined with the bioMérieux Advanced Expert System™ (AES) software using the interpretation standard CLSI M100-S22 (2012) and the interpretation guideline "Natural

- Resistance." For more information, please refer to Sanders, C. C. et al. "Potential Impact of the VITEK 2 System and the Advanced Expert System on the Clinical Laboratory of a University-Based Hospital." J. Clin. Microbiol. 39 (2001): 2379-2385. PubMed: 11427542.
- ⁴1 day at 37°C in an aerobic atmosphere on Mueller Hinton agar
- ⁵Minimum Inhibitory Concentration (MIC); MIC interpretation guidelines CLSI M100-S22 (2012)
- ⁶Rifampicin MIC interpretive standards are not available for *P. aeruginosa*. Strain Stone no. 130 is reported to be rifampicin-sensitive and does not grow in media containing rifampicin at a concentration of 100 µg/mL. For more information, please refer to Jacoby, G. A. "Properties of R plasmids Determining Gentamicin Resistance by Acetylation in *Pseudomonas aeruginosa*." Antimicrob. Agents Chemother. 6 (1974): 239-252. PubMed: 15830469.
- ⁷Streptomycin MIC interpretive standards are not available for *P. aeruginosa*. Strain Stone no. 130 contains IncP-2 plasmid pMG2, which confers resistance toward gentamicin, streptomycin and sulfonamides. Using an agar dilution method, the depositor determined that strains containing pMG2 have a MIC of 5000 µg/mL. Although strain Stone no. 130 can grow in the presence of a high level of streptomycin, BEI Resources did not confirm the presence of pMG2 by molecular methods. For more information, please refer to Jacoby, G. A. "Properties of R plasmids Determining Gentamicin Resistance by Acetylation in *Pseudomonas aeruginosa*." Antimicrob. Agents Chemother. 6 (1974): 239-252. PubMed: 15830469.
- ⁸Trimethoprim/sulfamethoxazole MIC interpretive standards are not available for *P. aeruginosa*; however most clinical isolates are resistant to trimethoprim/sulfamethoxazole. Strain Stone no. 130 contains IncP-2 plasmid pMG2, which confers resistance toward gentamicin, streptomycin and sulfonamides. Although strain Stone no. 130 can grow in the presence of a high level of trimethoprim/sulfamethoxazole, BEI Resources did not confirm the presence of pMG2 by molecular methods. For more information, please refer to Jacoby, G. A. "Properties of R plasmids Determining Gentamicin Resistance by Acetylation in *Pseudomonas aeruginosa*." Antimicrob. Agents Chemother. 6 (1974): 239-252. PubMed: 15830469 and Sanders, C. C. et al. "Potential Impact of the VITEK 2 System and the Advanced Expert System on the Clinical Laboratory of a University-Based Hospital." J. Clin. Microbiol. 39 (2001): 2379-2385. PubMed: 11427542.
- ⁹MIC result is based on the trimethoprim component of the test strip.
- ¹⁰Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, please refer to Auch, A.F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." Stand. Genomic Sci. 2 (2010): 117-134. PubMed: 21304684.
- ¹¹The whole genome of *P. aeruginosa*, strain Stone no. 130 (Contig Total Length ~ 6.3 megabase pairs) was sequenced using the Illumina® MiSeq® system and was assembled and analyzed with CLC Genomics Workbench Version 7.0.2.
- ¹²Purity of this lot was assessed for 7 days on Nutrient agar at 37°C in an aerobic atmosphere with 5% CO₂.

Figure 1: Colony Morphology



Date: 28 SEP 2017

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

