

## **Certificate of Analysis for NR-28525**

## **Human Respiratory Syncytial Virus, A2001/2-20**

Catalog No. NR-28525

**Product Description:** Cell lysate and supernatant from HEp-2 cells<sup>1</sup> infected with human respiratory syncytial virus, A2001/2-20

Lot<sup>2</sup>: 70005954 Manufacturing Date: 12JUN2017

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in HEp-2 Cells <sup>1</sup>	Cell rounding, detachment and syncytia formation	Cell rounding, detachment and syncytia formation
Sequencing of Species-Specific Region (733 nucleotides)	Consistent with human respiratory syncytial virus, A2001/2-20	100% sequence identity with human respiratory syncytial virus, A2001/2-20 (GenBank: JX069798)
Titer by TCID₅₀ Assay³,⁴ in HEp-2 Cells¹	Report results	2.8 × 10 <sup>5</sup> TCID <sub>50</sub> per mL
Sterility (21-day incubation) Harpo's HTYE broth <sup>5</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth	No growth
Mycoplasma Contamination  Agar and broth culture (14-day incubation at 37°C)  DNA Detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>&</sup>lt;sup>1</sup>HEp-2 cells: ATCC<sup>®</sup> CCL-23™

Date: 13 NOV 2017 Signatur

**BEI Resources Authentication** 

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

NR-28525\_70005954\_13NOV2017

<sup>&</sup>lt;sup>2</sup>Grown in Eagle's Minimum Essential Medium (ATCC® 30-2003™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 6 days at 37°C and 5% CO<sub>2</sub>

<sup>&</sup>lt;sup>3</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.
<sup>47</sup> days at 37°C and 5% CO<sub>2</sub>

<sup>&</sup>lt;sup>5</sup>Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.