

**Purified Zika Virus, PRVABC59**

**Catalog No. NR-50684**

**Product Description:** Purified virus from clarified cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells<sup>1</sup> infected with Zika virus (ZIKV), PRVABC59 (BEI Resources NR-50240)

**Passage History:** V3/V3 (Centers for Disease Control/BEI Resources); V# = Number of passages in Vero cells

**Lot<sup>2,3</sup>: 70006909**

**Manufacturing Date: 13SEP2017**

TEST	SPECIFICATIONS	RESULTS
<b>Infectivity in Vero E6 Cells<sup>1</sup></b>	Report results	Cell rounding and detachment
<b>Sequencing of Species-Specific Region (929 nucleotides)</b>	Consistent with ZIKV, PRVABC59	100% identity with ZIKV, PRVABC59 (GenBank: KU501215; KX087101)
<b>Titer by TCID<sub>50</sub> Assay in Vero E6 Cells<sup>1,4,5</sup></b>	Report results	9.3 × 10 <sup>5</sup> TCID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>Vero 76, clone E6: ATCC® CRL-1586™

<sup>2</sup>Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate (ATCC® 30-2003) and 2% fetal bovine serum (ATCC® 30-2020) for 9 days at 37°C with 5% CO<sub>2</sub>

<sup>3</sup>Purified from clarified cell lysate and supernatant by precipitation with PEG6000 and centrifugation through a 30% sucrose cushion

<sup>4</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>5</sup>7 days at 37°C with 5% CO<sub>2</sub>

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Date:** 17 JAN 2018

**Signature:** 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

