

***Anaplasma phagocytophilum*, Strain NCH-1**

Catalog No. NR-48807

For research use only. Not for human use.

Contributor:

Jason A. Carlyon, Ph.D., Department of Microbiology and Immunology, Virginia Commonwealth University, Richmond, Virginia, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Anaplasmataceae*, *Anaplasma*

Species: *Anaplasma phagocytophilum*

Strain: NCH-1

Original Source: *Anaplasma phagocytophilum* (*A. phagocytophilum*), strain NCH-1 was isolated in 1997 from the blood of a patient suffering from human granulocytic ehrlichiosis in Nantucket, Massachusetts, USA.^{1,2}

A. phagocytophilum is a Gram-negative, obligate intracytoplasmic bacteria that infects bone marrow-derived mammalian cells, predominantly of the myeloid lineage.³ The species was formerly known as *Ehrlichia phagocytophila* and classified in the family *Rickettsiaceae*, but subsequently reassigned to the family *Anaplasmataceae*, both families belonging to the order *Rickettsiales*.³ *A. phagocytophilum* is transmitted by *Ixodes persulcatus* complex ticks and is the causative agent of tick-borne fever of ruminants, equine granulocytic ehrlichiosis, and human granulocytic ehrlichiosis (HGE).^{3,4}

Material Provided:

Each vial contains approximately 1 mL of human promyelocytic leukemia cells (HL-60; ATCC® CCL-240™) infected with *A. phagocytophilum*, strain NCH-1, containing fetal bovine serum and DMSO (final concentrations are indicated on the Certificate of Analysis for each lot).

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-48807 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Host: HL-60 cells (ATCC® CCL-240™)

Growth Medium: RPMI-1640 medium containing 2 mM L-glutamine, 10 mM HEPES, 4500 mg/L glucose, 1 mM sodium pyruvate and 1500 mg/L sodium bicarbonate, supplemented with 10% fetal bovine serum, or equivalent.

Infection: Cells should be at a dilution of 1 × 10⁵ to 1 × 10⁶ cells/mL.

Incubation: 8 days at 37°C and 5% CO₂

Cytopathic Effect: Uninfected HL-60 cells are typically round with smooth borders. Infected cell borders are rough in appearance. It is recommended that replication of *A. phagocytophilum* be confirmed by IFA.

Citation:

Acknowledgment for publications should read “The following reagent was obtained through BEI Resources, NIAID, NIH: *Anaplasma phagocytophilum*, Strain NCH-1, NR-48807.”

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties.

Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Carlyon, J., Personal Communication.
2. Telford, S. R., et al. "Perpetuation of the Agent of Human Granulocytic Ehrlichiosis in a Deer Tick-Rodent Cycle." Proc. Natl. Acad. Sci. USA 93 (1996): 6209-6214. PubMed: 8650245.
3. Dumler, J. S., et al. "Reorganization of Genera in the Families *Rickettsiaceae* and *Anaplasmataceae* in the Order *Rickettsiales*: Unification of Some Species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, Descriptions of Six New Species Combinations and Designation of *Ehrlichia equi* and 'HGE agent' as Subjective Synonyms of *Ehrlichia phagocytophila*." Int. J. Syst. Evol. Microbiol. 51 (2001): 2145-2165. PubMed: 11760958.
4. Dumler, J. S., et al. "Human Granulocytic Anaplasmosis and *Anaplasma phagocytophilum*." Emerg. Infect. Dis. 11 (2005): 1828-1834. PubMed: 16485466.

ATCC® is a trademark of the American Type Culture Collection.

