

Mayaro Virus, Uruma

Catalog No. NR-49914

For research use only. Not for human use.

Contributor:

World Reference Center for Emerging Viruses and Arboviruses, University of Texas Medical Branch, Galveston, Texas, USA

Manufacturer:

BEI Resources

Product Description:

Virus Classification: *Togaviridae*, *Alphavirus*

Species: Mayaro virus

Strain/Isolate: Uruma

Original Source: Mayaro virus (MAYV), Uruma was isolated from a human in Uruma, Bolivia in March, 1955,¹ and contributed to WRCEVA by the Yale Arbovirus Research Unit, Rockefeller Funded Collection, Yale University, New Haven, Connecticut, USA. In order to remove contaminating mycoplasma, the first viral passage at BEI Resources was performed by lipofectamine-mediated transfection of extracted viral RNA.

MAYV is a New World alphavirus that is the etiologic agent of Mayaro fever, an acute febrile illness sometimes accompanied by severe and persistent arthritis. MAYV was first isolated in Trinidad in 1954, and there have been sporadic outbreaks of Mayaro fever in South America since. The enzootic transmission cycle of MAYV is not fully understood, but the occurrence of relatively large outbreaks of Mayaro fever³ and the competence of *Aedes* mosquitoes for transmission of MAYV⁴ suggest the potential for an urban human-mosquito-human transmission cycle to emerge.

There are two distinct genotypes of MAYV, D and L. Genotype D includes viruses isolated from all countries where MAYV has been detected, while genotype L strains have been found only in Brazil.^{1,2} Uruma is a D genotype virus.

Material Provided:

Each vial contains approximately 1 mL of cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells (Vero; ATCC® CCL-81™) infected with MAYV, Uruma.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-49914 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-

term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Host: *Cercopithecus aethiops* kidney epithelial cells (Vero; ATCC® CCL-81™)

Growth Medium: Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate supplemented with 2% fetal bovine serum, or equivalent

Infection: Cells should be 70% to 90% confluent

Incubation: 2 to 7 days at 37°C and 5% CO₂

Cytopathic Effect: Cell rounding and detachment

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH, as part of the WRCEVA program: Mayaro Virus, Uruma, NR-49914."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Powers, A. M., et al. "Genetic Relationships among Mayaro and Una Viruses Suggest Distinct Patterns of Transmission." Am. J. Trop. Med. Hyg. 75 (2006): 461-469. Pubmed: 16968922.
2. Auguste, A. J., et al. "Evolutionary and Ecological Characterization of Mayaro Virus Strains Isolated during an Outbreak, Venezuela, 2010." Emerg. Infect. Dis. 21 (2015): 1742-1750. Pubmed: 26401714.
3. LeDuc, J. W., F. Pinheiro, and A. Travassos da Rosa. "An Outbreak of Mayaro Virus Disease in Belterra, Brazil. II. Epidemiology." Am. J. Trop. Med. Hyg. 30 (1981): 682-688. Pubmed: 6266264.
4. Long, K. C., et al. "Experimental Transmission of Mayaro Virus by *Aedes aegypti*." Am. J. Trop. Med. Hyg. 85 (2011): 750-757. Pubmed: 21976583.

ATCC® is a trademark of the American Type Culture Collection.

