

Certificate of Analysis for NR-41795

Influenza B Virus, B/Florida/4/2006 (Yamagata Lineage)

Catalog No. NR-41795

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs¹ infected with influenza B virus, B/Florida/4/2006 (Yamagata Lineage)

Lot^{2,3}: 70010870 Manufacturing Date: 07DEC2017

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs ¹ Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin Coding Region (869 nucleotides)	Consistent with B/Florida/4/2006 (Yamagata Lineage)	100% identity with B/Florida/4/2006 (Yamagata Lineage) (GenBank: CY033876)
Titer by CEID ₅₀ Assay ^{4,5} in Embryonated Chicken Eggs ¹	Report results	1.6 × 10 ⁹ CEID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹⁰⁻ to 11-day-old SPF Embryonated Chicken Eggs acquired from Charles River, Norwich, Connecticut, USA

15 MAR 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

NR-41795 70010870 15MAR2018

²Derived from CDC ID No. 2007700596

³Grown in the allantoic cavity of embryonated chicken eggs¹ for 3 days at 35°C in a humidified chamber

⁴The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁵3 days at 35°C in a humidified chamber

⁶Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.