

Influenza B Virus, B/Sydney/507/2006 (Yamagata Lineage)

Catalog No. NR-36526

Product Description: Influenza B virus, B/Sydney/507/2006 (Yamagata Lineage) was isolated on June 16, 2006 from a human in Sydney, Australia. Each vial contains cell lysate and supernatant from Madin-Darby Canine Kidney (MDCK) cells¹ infected with influenza B virus, B/Sydney/507/2006 (Yamagata Lineage).

Lot²: 70012623

Manufacturing Date: 05FEB2018

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in MDCK Cells¹	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Hemagglutinin Coding Region (~ 970 nucleotides)	Consistent with B/Sydney/507/2006 (Yamagata Lineage)	100% identity with B/Sydney/507/2006 (Yamagata Lineage) (GenBank: EU124246.1)
Titer by TCID₅₀ Assay^{3,4} in MDCK Cells¹ by Cytopathic Effects	Report results	1.6 × 10 ⁸ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹MDCK; ATCC® CCL-34™

²Grown in Dulbecco's Modified Eagle Medium (ATCC® 30-2002™) supplemented with 0.125% bovine serum albumin (Invitrogen™ 15260-037) and 1 µg/mL L-1-tosylamido-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin for 3 days at 33°C and 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴8 days at 33°C and 5% CO₂

⁵Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

12 JUL 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.



BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370

Fax: 703-365-2898