

***Yersinia enterocolitica* subsp. *enterocolitica*, Strain 33114**

Catalog No. NR-214

(Derived from ATCC® 9610™)

For research use only. Not for human use.

Contributor:

ATCC®

Product Description:

Bacteria Classification: *Enterobacteriaceae*, *Yersinia*

Agent: *Yersinia enterocolitica* subsp. *enterocolitica*^{1,2}

Biotype/Biovar: 1

Serotype/Serovar: 0:8

Phagovar: X_z

Type Strain: 33114

Original Source:^{3,4} Isolated from facial abscesses of an adult human with a chronic, glanders-like infection of the face in New York, 1934

Comments: *Yersinia enterocolitica* subsp. *enterocolitica* (*Y. enterocolitica* subsp. *enterocolitica*), strain 33114 was deposited at ATCC® in 1944 by Julia M. Coffee, Associate Bacteriologist, New York Department of Health, Division of Laboratories and Research, Albany, New York.

Y. enterocolitica subsp. *enterocolitica* is a significant food-borne enteropathogen which causes gastroenteritis. *Y. enterocolitica* subsp. *enterocolitica* is an extremely heterogeneous species, encompassing six biotypes and currently more than 50 serogroups, not all of which can cause disease.⁵ It is of particular concern to the food industry because it is a psychrotrophic pathogen able to proliferate at temperatures approaching 0°C.

Y. enterocolitica subsp. *enterocolitica* is a non-spore-forming, gram-negative, rod-shaped coccobacillus. Virulence-associated genes are located on the chromosome and on the pYV (~ 64 to 75 kb) plasmid found in typical virulent strains of *Y. enterocolitica* subsp. *enterocolitica*.⁶ This plasmid encodes a type III secretion system involved in the delivery of virulence proteins that contribute to internalization into host cells.⁷

The presence of the pYV plasmid in NR-214 has been confirmed by gel electrophoresis of extracted DNA.

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in 0.5X Tryptic Soy Broth supplemented with 10% glycerol.

Packaging/Storage:

NR-214 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen

freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Brain Heart Infusion Broth or Tryptic Soy Broth

Tryptic Soy Agar or Sheep Blood Agar

Incubation:

Temperature:⁸ 26°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; thaw slowly.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 26°C for 24–48 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Yersinia enterocolitica* subsp. *enterocolitica*, Strain 33114, NR-214."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 4th ed. Washington, DC: U.S. Government Printing Office, 1999. HHS Publication No. (CDC) 93-8395. This text is available online at www.cdc.gov/od/ohs/biosfty/bmbl4/bmbl4toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this

product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC[®], their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. "Validation List No. 75." Int. J. Syst. Evol. Microbiol. 50 (2000): 1415–1417.
2. Neubauer, H., et al. "*Yersinia enterocolitica* 16S rRNA Gene Types Belong to the Same Genospecies but Form Three Homology Groups." Int. J. Med. Microbiol. 290 (2000): 61–64. PubMed: 11043982.
3. McIver, M. A. and R. M. Pike. "Chronic Glanders-Like Infection of Face Caused by an Organism Resembling *Flavobacterium pseudomallei* Whitmore." Clinical Misc. Mary Imogene Basset Hosp. 1 (1934): 16–21.
4. Schleifstein, J. and M. B. Coleman. "Unidentified Microorganisms Resembling *B. lignieri* and *Past. pseudotuberculosis* Pathogenic for Man." N.Y. State J. Med. 39 (1939): 1749–1753.
5. Viridi, J. S. and P. Sachdeva. "Molecular Heterogeneity in *Yersinia enterocolitica* and 'Y. enterocolitica-Like' Species – Implications for Epidemiology, Typing and Taxonomy." FEMS Immunol. Med. Microbiol. 45 (2005): 1–10. PubMed: 15985218.
6. Bottone, E. J. "*Yersinia enterocolitica*: The Charisma Continues." Clin. Microbiol. Rev. 10 (1997): 257–276. PubMed: 9105754.
7. Snellings, N. J., M. Popek, and L. E. Lindler. "Complete DNA Sequence of *Yersinia enterocolitica* Serotype O:8 Low-Calcium-Response Plasmid Reveals a New Virulence Plasmid-Associated Replicon." Infect. Immun. 69 (2001): 4627–4638. PubMed: 11402007.
8. Chu, M. C. Laboratory Manual of Plague Diagnostic Tests. Centers for Disease Control and Prevention, Atlanta, 2000.
9. Park, S., L. T. Smith, and G. M. Smith. "Role of Glycine Betaine and Related Osmolytes in Osmotic Stress Adaptation in *Yersinia enterocolitica* ATCC 9610." Appl. Environ. Microbiol. 61 (1995): 4378–4381. PubMed: 16535192.

ATCC[®] is a trademark of the American Type Culture Collection.