

Influenza A Virus, A/Puerto Rico/8/1934, BPL-Inactivated

Catalog No. NR-19325

Product Description: NR-19325 is a preparation of influenza A virus, A/Puerto Rico/8/1934 that has been inactivated with beta-propiolactone (BPL). The source virus used for the inactivation was BEI Resources NR-348, Lot No. 4729801.

Lot: 59300930

Manufacturing Date: 28JUN2010

TEST	SPECIFICATIONS	RESULTS
Concentration (in CEID₅₀ equivalents)	Report results	3.5 × 10 ⁶ CEID ₅₀ equivalents per mL
Innocuity Test (Screening for Viral Inactivation in Eggs)^{1,2} NR-19325, Influenza A Virus, A/Puerto Rico/8/1934, BPL-inactivated ³ 1 st round of amplification (1:10) 2 nd round of amplification (neat) 3 rd round of amplification (neat) NR-348, Influenza A Virus, A/Puerto Rico/8/1934 Positive Control 1 st round of amplification (1:10) 2 nd round of amplification (1:10) 3 rd round of amplification (1:10)	No HA activity detected No HA activity detected No HA activity detected HA activity detected HA activity detected HA activity detected	No HA activity detected No HA activity detected No HA activity detected HA activity detected HA activity detected HA activity detected
Functional Activity RNA detection by PCR of extracted Test Article nucleic acid Influenza A virus primer and probe set Influenza H1 seasonal primer and probe set Influenza H3 primer and probe set Influenza H5a primer and probe set Influenza H5b primer and probe set Influenza B virus primer and probe set Antigenicity using BD™ Directigen™ EZ Flu A+B (Cat. No. 256050) Influenza A Influenza B	Detected Detected None detected None detected None detected None detected Report results Report results	Detected Detected None detected None detected None detected None detected Reactive Non-reactive
Sterility (21-day incubation) Harpo's HTYE broth ³ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected	No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected No growth detected
Mycoplasma Contamination DNA detection by PCR of extracted Test Article nucleic acid	None detected	None detected

¹9- to 11-day old embryonated chicken eggs were inoculated with 0.2 mL of the indicated test sample and incubated at 35°C for 2 days. Allantoic fluid from the first round of amplification was tested for HA activity and 0.2 mL was inoculated into 9- to 11-day old embryonated chicken eggs and incubated at 35°C for 2 days. Allantoic fluid from the second round of amplification was tested for HA activity and 0.2 mL was inoculated into 9- to 11-day old embryonated chicken eggs and incubated at 35°C for 2 days. Allantoic fluid from the third round of amplification was tested for HA activity.

²The inoculum for the first round of amplification for the Positive Control (NR-348) was 3.2×10^7 CEID₅₀ per egg. The inoculum for the first round of amplification for the BPL-inactivated Test Article (NR-19325) was 2.8×10^7 CEID₅₀ equivalents per egg.

³Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

27 NOV 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

