

***Toxoplasma gondii*, ME49 (B7 clone)**

Catalog No. NR-225

(Derived from ATCC® 50840™)

For research use only. Not for human use.

Contributor:

L. David Sibley, Professor, Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri.

Product Description:

Protozoa Classification: *Apicomplexa*, *Toxoplasma*

Species: *Toxoplasma gondii*

Strain: ME49 (B7 clone)^{1,2}; also referred to as P and PLK strain

Original Source: *Toxoplasma gondii*, ME49 (originally isolated from a sheep in California³ and available as BEI Resources NR-362) was passed singly through a cat to obtain a line that was capable of producing oocysts, and then further cloned by limiting dilution to produce the B7 clone.^{1,2}

Comment: The ME49 (B7 clone) strain was deposited at ATCC®, in 2000 by Dr. L. David Sibley of the Washington University School of Medicine in St. Louis, Missouri, as Type II genotype: SAG1-1; SAG2-2; SAG3-2. Extensive testing of ME49, ME49 (B7 clone) and PTG (an independent clone of ME49) indicated that ME49 was effectively already a biological clone.¹

Toxoplasma gondii, ME49 (B7 clone) is the Type II parental strain that was used in a genetic cross with the genotype Type III parental strain CTG ARA-SNF (also known as CTG ARA-A^R/SNF^R and CEP ARA-A^R/SNF^R and available as BEI Resources NR-10151) to test the feasibility of inter-strain crosses.^{1,4,5} Strain CTG ARA-SNF is the product of a genetic cross between singly resistant parental clones of the CTG (CEP) strain obtained by chemical mutagenesis.⁶

Toxoplasma gondii (*T. gondii*) is an obligate intracellular protozoan parasite of the phylum *Apicomplexa* that is the causal agent of toxoplasmosis. *T. gondii* has a highly unusual, clonal population structure comprised of three widespread genotypes referred to as type I, type II, and type III. Type I are the highly virulent, type II are nonvirulent and give rise to chronic infections in humans, and type III is also nonvirulent but mainly associated with chronic infections in animals.^{7,8} The genetic basis for the differences between the three lineages has been recently examined by mapping virulence in F1 progeny derived from crosses between the different *T. gondii* lineages.^{9,10}

T. gondii ME49 (B7 clone) is commonly associated with AIDS. The whole genome shotgun sequence of ME49 (B7 clone) has been deposited in GenBank (ABPA01000000).

Note: This strain is also available as NR-10150 which has had additional characterization tests performed.

Material Provided:

Each vial of NR-225 contains approximately 0.5 mL of culture in cryopreservative. Please see Appendix I below for cryopreservation instructions.

Packaging/Storage:

NR-225 was packaged aseptically in screw-capped plastic cryovials and is provided frozen on dry ice. The product should be stored at cryogenic temperature (-130°C or colder), preferably in the vapor phase of a liquid nitrogen freezer. If liquid nitrogen storage facilities are not available, frozen cryovials may be stored at -70°C or colder for approximately one week. Note: Do not under any circumstances store vials at temperatures warmer than -70°C. Storage under these conditions will result in the death of the culture.

To insure the highest level of viability, the culture should be initiated immediately upon receipt. Any warming of the product during shipping and transfer must be avoided, as this will adversely affect the viability of the product. For transfer between freezers and for shipping, the product may be placed on dry ice for brief periods, although use of a portable liquid nitrogen carrier is preferred. Please read the following recommendations prior to using this material.

Growth Conditions:

ATCC medium 2222: Cell cultivation medium for parasites (Dulbecco's Minimal Essential Medium), adjusted to contain 10% (v/v) heat-inactivated fetal bovine serum (HIFBS)

Human foreskin fibroblast cells (ATCC® CRL-1634™)

Incubation:

Temperature: 35 to 37°C

Atmosphere: 95% air, 5% CO₂

Propagation:

1. To establish a culture from the frozen state, place a vial in a 35 to 37°C water bath. Thawing time is approximately 2 to 3 minutes. Do not agitate the vial. Do not leave the vial in the water bath after it is thawed.
2. Immediately after thawing, aseptically transfer the contents to a tissue culture flask containing a fresh monolayer of human foreskin fibroblast cells (ATCC® CRL-1634™) and 10 mL of ATCC medium 2222 containing 10% (v/v) HIFBS.
3. Outgas the flask for 10 seconds with a 95% air, 5% CO₂ gas mixture.
4. Incubate in a 35 to 37°C CO₂ incubator with the caps screwed on tightly. Observe the culture daily under an inverted microscope for the presence of parasitophorous vacuoles.

Maintenance:

1. Remove the medium from a fresh confluent monolayer of human foreskin fibroblast cells in a tissue culture flask

and replace it with 10 mL medium containing 10% (v/v) HIFBS.

- Remove the medium from the *Toxoplasma* culture when approximately 50% of the human foreskin fibroblast cell monolayer has lysed. Centrifuge the parasites that had been released into the medium at 1300 x g for 10 minutes.
- Remove the supernatant and resuspend the cell pellet in a small volume (0.5 to 1.0 mL) of ATCC medium 2222 or phosphate buffered saline (PBS). Transfer the resuspended pellet to the fresh flask of human foreskin fibroblast cells, prepared in step 1 above. Follow steps 3 and 4 in Propagation.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Toxoplasma gondii*, ME49 (B7 Clone), NR-225."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmb15/bmb15toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

- Ajioaka, J. W. and L. D. Sibley. "Development and Application of Classical Genetics in *Toxoplasma gondii*." *Toxoplasma gondii: The Model Apicomplexan. Perspectives and Methods*. Ed. L. M. Weiss and K. Kim. London: Academic Press, 2007. 374-375.
- Kasper, L. H. and P. L. Ware. "Recognition and Characterization of Stage-Specific Oocyst/Sporozoite Antigens of *Toxoplasma gondii* by Human Antisera." *J. Clin. Invest.* 75 (1985): 1570-1577. PubMed: 2581998.
- Lunde, M. N. and L. Jacobs. "Antigenic Differences between Endozoites and Cystozoites of *Toxoplasma gondii*." *J. Parasitol.* 69 (1983): 806-808. PubMed: 6200590.
- Sibley, L. D., et al. "Generation of a Restriction Fragment Length Polymorphism Linkage Map for *Toxoplasma gondii*." *Genetics* 132 (1992): 1003-1015. PubMed: 1360931.
- Khan, A., et al. "Composite Genome Map and Recombination Parameters Derived from Three Archetypal Lineages of *Toxoplasma gondii*." *Nucleic Acids Res.* 33: 2980-2992. PubMed: 15911631.
- Pfefferkorn, E.R. and L. H. Kasper. "*Toxoplasma gondii*: Genetic Crosses Reveal Phenotypic Suppression of Hydroxyurea Resistance by Fluorodeoxyuridine Resistance." *Exp. Parasitol.* 55 (1983): 207-218. PubMed: 6219892.
- Sibley, L. D. and J. C. Boothroyd. "Virulent Strains of *Toxoplasma gondii* Comprise a Single Clonal Lineage." *Nature* 359 (1992): 82-85. PubMed: 1355855.
- Howe, D. K. and L. D. Sibley. "*Toxoplasma gondii* Comprises Three Clonal Lineages: Correlation of Parasite Genotype with Human Disease." *J. Infect. Dis.* 172 (1995): 1561-1566. PubMed: 7594717.
- Saeij, J. P., et al. "Polymorphic Secreted Kinases are Key Virulence Factors in Toxoplasmosis." *Science* 314 (2006): 1780-1783. PubMed: 17170306.
- Taylor, S., et al. "A Secreted Serine-Threonine Kinase Determines Virulence in the Eukaryotic Pathogen *Toxoplasma gondii*." *Science* 314 (2006): 1776-1780. PubMed: 17170305.

ATCC® is a trademark of the American Type Culture Collection.



APPENDIX I: CRYOPRESERVATION

1. To harvest the *Toxoplasma* culture, detach any remaining tissue culture cells (infected and uninfected) by scraping the surface of the flask with a cell scraper.
2. Transfer the cell suspension (including parasites) to 15 mL plastic centrifuge tubes. Centrifuge at 1300 x g for 10 min.
3. Remove all but 0.5 mL of the supernatant from each tube, resuspend the cell pellets, and pool them to a single tube.
4. Pass the resulting cell suspension through a syringe equipped with a 27-gauge ½-inch needle to break up any remaining cells.
5. Adjust the parasite concentration to 2.0 to 4.0 x 10⁷ cells/mL with fresh medium [Cell cultivation medium for parasites (ATCC® medium 2222) or Dulbecco's PBS (ATCC® 30-2200) can be used].
Note: If the concentration of parasites is too low, centrifuge at 1300 x g for 10 min and resuspend in a smaller volume of fresh medium to yield the desired parasite concentration.
6. Mix equal volumes of parasite suspension and fresh medium or PBS containing 20% DMSO and 50% HIFBS to yield a final concentration of 1.0 to 2.0 x 10⁷ cells/mL in 10% DMSO, 25% HIFBS. The freezing process should start 15 to 30 minutes following the addition of cryoprotective solution to the parasite suspension.
Note: To prevent culture contamination, penicillin-streptomycin solution (ATCC® 30-2300) may be added to a final concentration of 50 to 100 U/mL penicillin and 50 to 100 µg/mL streptomycin.
7. Dispense 0.5 mL aliquots into 1 to 2 mL sterile plastic screw-capped vials for cryopreservation.
8. Place the vials in a controlled rate freezing unit. From room temperature cool the vials at -1°C/min to -40°C. If the freezing unit can compensate for the heat of fusion, maintain rate at -1°C/min through this phase. At -40°C, plunge vials into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing container. Place the container at -80°C for 1.5 to 2 hours and then plunge vials into liquid nitrogen.
9. Store in either the vapor or liquid phase of a nitrogen refrigerator (-130°C or colder).