

**SARS-Related Coronavirus 2, Isolate USA-WA1/2020, Heat Inactivated**

**Catalog No. NR-52286**

This reagent is the tangible property of the U.S. Government.

**Product Description:**

NR-52286 is a preparation of SARS-Related Coronavirus 2 (SARS-CoV-2), isolate USA-WA1/2020 that has been inactivated by heating to 65°C for 30 minutes.

**Lot: 70037779**

**Manufacturing Date: 09APR2020**

TEST	SPECIFICATIONS	RESULTS
<b>Pre-Inactivation Titer by TCID<sub>50</sub> Assay in Vero E6 Cells<sup>1</sup></b> (7 days at 37°C and 5% CO <sub>2</sub> )	Report results	1.6 × 10 <sup>5</sup> TCID <sub>50</sub> per mL
<b>Pre-Inactivation Sterility</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>2</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Pre-Inactivation Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected
<b>Genome Copy Number Using BioRad QX200 Droplet Digital PCR (ddPCR™) System<sup>3</sup></b> (Post vial; 9 replicates)	Report results	1.77 × 10 <sup>8</sup> genome equivalents/mL
<b>Virus Inactivation</b> 10% of total bulk heat-treated preparation inoculated on Vero E6 cells and evaluated for cytopathic effect and presence of viral RNA by qRT-PCR <sup>4</sup>	No viable virus detected	No viable virus detected

<sup>1</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>2</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

<sup>3</sup>The GE value reported is obtained using Qiagen RNA extraction kit (Cat 52904).

<sup>4</sup>The inactivated virus preparation was plated on Vero E6 cells and incubated for 14 days at 37°C and 5% CO<sub>2</sub>; cell lysate and supernatant from these cultures were blind passaged on fresh monolayers of Vero E6 cells and again incubated for 14 days at 37°C and 5% CO<sub>2</sub>.

/Heather Couch/

Heather Couch

06 AUG 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

