

SARS-Related Coronavirus 2, Isolate USA-CA4/2020

Catalog No. NR-52386

Product Description:

Severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), isolate USA-CA4/2020 was isolated from an oropharyngeal swab from a human patient in California, USA on January 29, 2020. NR-52386 lot 70034885 was produced by infecting *Cercopithecus aethiops* kidney cells (Vero E6; ATCC® CRL-1586™) with the deposited material in Eagle's Minimum Essential Medium (ATCC 30-2003™) supplemented with 2% fetal bovine serum (ATCC 30-2020™) for 4 days at 37°C with 5% CO₂.

Passage History:

V(2)/VE6(2) (Centers for Disease Control and Prevention/BEI Resources); V = Vero cells, VE6 = Vero E6 cells

Lot: 70034885

Manufacturing Date: 18MAY2020

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero E6 Cells	Cell rounding and detachment	Cell rounding and detachment
Next-Generation Sequencing (NGS) of Complete Genome Using Illumina® iSeq™ 100 Platform (Refer to Appendix I for NGS information)	≥ 98% identity with SARS-CoV-2, isolate USA-CA4/2020 (GenBank: MT027063.1)	99.99% identity with SARS-CoV-2, isolate USA-CA4/2020 (GenBank: MT027063.1)
Titer by TCID₅₀ Assay in Vero E6 Cells by Cytopathic Effect¹ (9 days at 37°C and 5% CO ₂)	Report results	1.6 × 10 ⁶ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ² Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

²Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

14 AUG 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.



APPENDIX I: NGS Information for NR-52386 lot 70034885

Sequence analysis resulted in the discovery of three SNPs when compared to GenBank MT027063.1 (see Table below). Quality scores over 60 indicate it is improbable that the variant call is incorrect.

Position in NR-52386_70034885 Sequence	Position in MT027063.1	Reported MT027063.1 Sequence	Identified Alternative Base	Quality	Variant Type	Length of Variant	Frequency of Variant
22195	22206	A	G	53	SNP	1	0.1889764
28596	23607	G	T	221	SNP	1	0.8170213
26187	26198	C	T	221	SNP	1	0.7892562