

Vector pET-11a Containing the SARS-Related Coronavirus 2, Wuhan-Hu-1 Nucleocapsid Protein C-Terminal Domain Gene

Catalog No. NR-52434

This reagent is the tangible property of the U.S. Government.

Product Description:

The C-terminal domain (CTD) of the nucleocapsid (N) protein (amino acids 247 to 364; GenPept: [QHD43423](#)) from severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), Wuhan-Hu-1 (GenBank: [MN908947](#)) was codon optimized, tagged with a tobacco etch virus (TEV) cleavable N-terminal hexa-histidine tag and cloned into the pET-11a plasmid. The beta-lactamase gene, *bla*, provides transformant selection through ampicillin resistance in *Escherichia coli* (*E. coli*). The deposited plasmid was transformed into One Shot™ TOP10 *Escherichia coli* (Invitrogen™ C404003), grown in Luria-Bertani broth with ampicillin (50 µg per mL) for 1 day at 37°C in an aerobic atmosphere, extracted using a Plasmid Plus Maxi Kit (QIAGEN® 12963) and vialled in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0).

Lot: 70035138

Manufacturing Date: 06MAY2020

TEST	SPECIFICATIONS	RESULTS
Next-Generation DNA Sequencing	Report results	6100 base pairs ¹
Genotypic Analysis Sequencing of N CTD insert (~460 base pairs) N-terminal His ₆ tag N-terminal TEV protease site	100% sequence identity to depositor's sequence His ₆ tag sequence confirmed TEV protease site sequence confirmed	100% sequence identity to depositor's sequence ² His ₆ tag sequence confirmed TEV protease site sequence confirmed
Antibiotic Resistance Ampicillin (encoded by beta-lactamase gene <i>bla</i>) ³	<i>bla</i> sequence present	<i>bla</i> sequence present
Concentration by PicoGreen® Measurement	≥ 2 µg/mL	0.09 µg in 20 µL per vial (5 µg/mL)
Amount per Vial	Report results	0.09 µg per vial
OD₂₆₀/OD₂₈₀ Ratio (pre-vial)	1.7 to 2.1	2.0
Effective Bacterial Transformation Invitrogen™ One Shot™ TOP10 <i>Escherichia coli</i>	≥ 50 colonies per ng	> 500 colonies per ng

¹The sequence was assembled pre-vial using the depositor's predicted sequence as the reference sequence. The complete plasmid sequence and map are provided on the BEI Resources webpage.

²The NR-52434 insert was codon optimized but is consistent with the SARS-CoV-2, Wuhan-Hu-1 N CTD protein (amino acids 247 to 364; GenPept: QHD43423).

³The antibiotic ampicillin degrades quickly during growth. Bacterial stationary phase should be minimized during plasmid expansion to avoid plasmid loss and increased antibiotic concentrations may be necessary.

/Heather Couch/

Heather Couch

26 MAY 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by ATCC® and the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

