

***Mycobacterium tuberculosis* subsp. *tuberculosis*, Strain H37Ra**

Catalog No. NR-122

(Derived from ATCC® 25177™)

For research use only. Not for use in humans.

Contributor:

ATCC®

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Mycobacteriaceae*, *Mycobacterium*

Species: *Mycobacterium tuberculosis* subsp. *tuberculosis*¹

Strain: H37Ra

Source: *Mycobacterium tuberculosis* (*M. tuberculosis*) subsp. *tuberculosis*, strain H37Ra is an attenuated strain derived from the virulent parent strain H37. Strain H37 was isolated in 1905 from the sputum of a patient with chronic pulmonary tuberculosis.²

Comment: *M. tuberculosis* subsp. *tuberculosis*, strain H37Ra was deposited to ATCC® in 1969 by Carl L. Larson, M.D., Stella Duncan Institute, University of Montana, Missoula, Montana, USA. The complete genome of *M. tuberculosis*, strain H37Ra has been sequenced (GenBank: [CP000611](https://www.ncbi.nlm.nih.gov/nuclseq/CP000611)).³

M. tuberculosis subsp. *tuberculosis* is an acid-fast, Gram-positive, non-motile, rod-shaped aerobic bacterium. It is the causative agent of tuberculosis (TB) and is responsible for more morbidity in humans than any other bacterial disease. *M. tuberculosis* subsp. *tuberculosis* is a slow-growing pathogen with a thick, lipid-rich cell wall, lending bacilli the unusual propensity to shut down its metabolism in the face of adverse conditions and enter a latent phase in which it displays phenotypic resistance to antibiotic therapy. The primary focus of infection is the lungs, with TB being spread by infectious aerosols produced by coughing. The spread of multidrug-resistant (MDR) and extensively drug-resistant (XDR) TB is a major medical and public health concern.^{4,5,6,7,8,9}

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Middlebrook 7H9 broth with Middlebrook ADC Enrichment supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-122 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Middlebrook 7H9 broth with Middlebrook ADC Enrichment
Middlebrook 7H10 agar with Middlebrook OADC Enrichment

Incubation:

Temperature: 37°C
Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate at 37°C for 2 to 6 weeks.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Mycobacterium tuberculosis* subsp. *tuberculosis*, Strain H37Ra, NR-122."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale. This material may be subject to third party patent rights.

References:

1. "Notification that New Names and New Combinations Have Appeared in Volume 49, Part 3, of the IJSB." Int. J. Syst. Bacteriol. 49 (1999): 1327-1328.
2. Steenken, W., Jr., W. H. Oatway, Jr. and S. A. Petroff. "Biological Studies of the Tubercle Bacillus: III. Dissociation and Pathogenicity of the R and S Variants of the Human Tubercle Bacillus (H37)." J. Exp. Med. 60 (1934): 515-540.
3. Zheng, H., et al. "Genetic Basis of Virulence Attenuation Revealed by Comparative Genomic Analysis of *Mycobacterium tuberculosis* Strain H37Ra versus H37Rv." PLoS One 3 (2008): e2375. PubMed: 18584054.
4. Dye, C. "Doomsday Postponed? Preventing and Reversing Epidemics of Drug-resistant Tuberculosis." Nat. Rev. Microbiol. 7 (2009): 81-87. PubMed: 19079354.
5. Chan, E. D. and M. D. Iseman. "Multidrug-resistant and Extensively Drug-resistant Tuberculosis: A Review." Curr. Opin. Infect. Dis. 21 (2008): 587-595. PubMed: 18978526.
6. Balganes, T. S., P. M. Alzari and S. T. Cole. "Rising Standards for Tuberculosis Drug Development." Trends Pharmacol. Sci. 29 (2008): 576-581. PubMed: 18799223.
7. Grandjean, L. and D. A. J. Moore. "Tuberculosis in the Developing World: Recent Advances in Diagnosis with Special Consideration of Extensively Drug-Resistant Tuberculosis." Curr. Opin. Infect. Dis. 21 (2008): 454-461. PubMed: 18725793.
8. Murphy, D. J. and J. R. Brown. "Novel Drug Target Strategies against *Mycobacterium tuberculosis*." Curr. Opin. Microbiol. 11 (2008): 422-427. PubMed: 18801459.
9. Hoft, D. F. "Tuberculosis Vaccine Development: Goals, Immunological Design, and Evaluation." Lancet 372 (2008): 164-175. PubMed: 18620952.
10. Cubillos-Ruiz, A., J. Morales and M. M. Zambrano. "Analysis of the Genetic Variation in *Mycobacterium tuberculosis* Strains by Multiple Genome Alignments." BMC Res. Notes 1 (2008): 110. PubMed: 18992142.
11. Adams, K. L., et al. "Reagentless Detection of *Mycobacteria tuberculosis* H37Ra in Respiratory Effluents in Minutes." Anal. Chem. 80 (2008): 5350-5357. PubMed: 18558726.
12. Kelly, D. M., et al. "Bystander Macrophage Apoptosis after *Mycobacterium tuberculosis* H37Ra Infection." Infect. Immun. 76 (2008): 351-360. PubMed: 17954721.

13. Sharma, D. and J. S. Tyagi. "The Value of Comparative Genomics in Understanding Mycobacterial Virulence: *Mycobacterium tuberculosis* H37Ra Genome Sequencing – A Worthwhile Endeavor." J. Biosci. 32 (2007): 185-189. PubMed: 17435310.
14. Chen, M., H. Gan and H. G. Remold. "A Mechanism of Virulence: Virulent *Mycobacterium tuberculosis* Strain H37Rv, but not Attenuated H37Ra, Causes Significant Mitochondrial Inner Membrane Disruption in Macrophages Leading to Necrosis." J. Immunol. 176 (2006): 3707-3716. PubMed: 16517739.
15. Soto, C. Y., et al. "Simple and Rapid Differentiation of *Mycobacterium tuberculosis* H37Ra from *M. tuberculosis* Clinical Isolates through Two Cytochemical Tests Using Neutral Red and Nile Blue Stains." J. Clin. Microbiol. 40 (2002): 3021-3024. PubMed: 12149369.
16. Acharya, P. V. and D. S. Goldman. "Chemical Composition of the Cell Wall of the H37Ra Strain of *Mycobacterium tuberculosis*." J. Bacteriol. 102 (1970): 733-739. PubMed: 4988039.

ATCC® is a trademark of the American Type Culture Collection.

