



## NIH AIDS Reagent Program

20301 Century Boulevard  
Building 6, Suite 200  
Germantown, MD 20874  
USA

Phone: 240 686 4740  
Fax: 301 515 4015  
aidsreagent.org

### DATA SHEET

<b>Reagent:</b>	☒ HTLV-1 OS-P2 Infected Cells
<b>Catalog Number:</b>	1393
<b>Lot Number:</b>	98088
<b>Release Category:</b>	A
<b>Provided:</b>	1 vial frozen cells, $4 \times 10^6$ cells/ml (90% viability).
<b>Cell Type:</b>	Subclone of HTLV-I infected CD8+ human T-lymphocyte cell line. Morphology is similar to T-cell lines. Syncytium formation is rare.
<b>Propagation Medium:</b>	RPMI 1640, 85%; fetal bovine serum, 15%.
<b>Freeze Medium:</b>	RPMI 1640, 82.5%; fetal bovine serum, 10%; DMSO, 7.5%.
<b>Growth Characteristics:</b>	Thaw rapidly and slowly dilute the cells with 37°C medium. Initiate culture at $5 \times 10^5$ cells/mL. Replace 80% of the medium every 3-4 days, and split the cells approximately 1:2 every 6-8 days or as necessary. Do not maintain the cells at concentrations greater than $2.0 \times 10^6$ cells/mL. Approximate doubling time is every six days. These cells grow clumped in suspension. The culture medium can be supplemented with low concentrations of IL-2 to enhance growth.
<b>Sterility:</b>	Negative for bacteria, fungi, and mycoplasma.
<b>Description:</b>	HTLV-1 OS-P2 infected cells secrete lymphotoxin, several other cytokines (e.g., GM-CSF), abundant HTLV-I virus particles and HTLV-I antigen (e.g., viral p24).
<b>Special Characteristics:</b>	This cell line was obtained by culturing normal human PBMCs with cerebrospinal fluid from an HTLV-I-seropositive patient with HTLV-I-associated myelopathy/tropical spastic paraparesis (HAM/TSP). Moderately infectious when inoculated into rabbits. HTLV-1 OS-P2 infected cells are CD4+ and maintains many normal cell surface markers (e.g., CD3, CD2).

---

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

**Recommended Storage:** Liquid nitrogen.

**Contributor:** Dr. Michael D. Lairmore.

**References:** Lairmore MD, Roberts B, Frank D, Rovnak J, Weiser MG, Cockerel, GL. Comparative biological responses of rabbits infected with human T-lymphotropic virus type I isolates from patients with lymphoproliferative and neurodegenerative disease. *Int J Cancer* **50**:124-130, 1992.

McKendall RR, Oas J, Lairmore MD. HTLV-I-associated myelopathy endemic in Texas-born residents and isolation of virus from CSF cells. *Neurology* **41**:831-836, 1991.

**NOTE:** Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HTLV-1 OS-P2 Infected Cells from Dr. Michael D. Lairmore." Please include the references cited above in any publications.

**Last Updated** July 02, 2018

---

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.