

NIH AIDS Reagent Program

20301 Century Boulevard Building 6, Suite 200 Germantown, MD 20874 USA

Phone: 240 686 4740 Fax: 301 515 4015 aidsreagent.org

DATA SHEET

Reagent:	Anti-HIV-1 gp120 Monoclonal (2G12)
Catalog Number:	1476
Lot Number:	140426
Release Category:	A
Provided:	250 μg of purified antibody at 1 mg/mL in 2 mM acetic acid, 10% maltose (sterile and does not contain preservatives)
Description:	A recombinant monoclonal antibody to HIV-1 gp120
Host:	Human
Titer:	The user should determine the optimal concentration for any application.
Special Characteristics:	This antibody was produced in a recombinant CHO cell expression system and purified by protein A affinity chromatography. This antibody originates from a HIV-1 positive human donor. Please see the <u>LANL HIV Molecular Database</u> for more information.
	This antibody neutralizes a broad variety of SHIV variants and HIV-1 laboratory strains and primary isolates. The epitope is conformational and carbohydrate-dependent. It is directed against N-linked glycans in the C2, C3, V4, and C4 domains of gp120.
Recommended Storage:	Keep the reagent at 4°C for short term storage and at -80°C for long term storage. Avoid freeze-thaw cycles as reagent degradation may result.
Contributor:	DAIDS, NIAID (Produced by Polymun Scientific)

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS. Buchacher, A., Predl, R., Strutzenberger, K., Steinfellner, W., Trkola, A., Purtscher, M., . . . et al. (1994). Generation of human monoclonal antibodies against HIV-1 proteins; electrofusion and Epstein-Barr virus transformation for peripheral blood lymphocyte immortalization. AIDS Res Hum Retroviruses, 10(4), 359-369. doi:10.1089/aid.1994.10.359 PUBMED

Crawford, J. M., Earl, P. L., Moss, B., Reimann, K. A., Wyand, M. S., Manson, K. H., . . . Montefiori, D. C. (1999). Characterization of primary isolate-like variants of simian-human immunodeficiency virus. J Virol, 73(12), 10199-10207. <u>PUBMED</u>

Etemad-Moghadam, B., Sun, Y., Nicholson, E. K., Karlsson, G. B., Schenten, D., & Sodroski, J. (1999). Determinants of neutralization resistance in the envelope glycoproteins of a simian-human immunodeficiency virus passaged in vivo. J Virol, 73(10), 8873-8879. <u>PUBMED</u>

Mascola, J. R., Lewis, M. G., Stiegler, G., Harris, D., VanCott, T. C., Hayes, D., . . . Birx, D. L. (1999). Protection of Macaques against pathogenic simian/human immunodeficiency virus 89.6PD by passive transfer of neutralizing antibodies. J Virol, 73(5), 4009-4018. <u>PUBMED</u>

Trkola, A., Purtscher, M., Muster, T., Ballaun, C., Buchacher, A., Sullivan, N., . . . Katinger, H. (1996). Human monoclonal antibody 2G12 defines a distinctive neutralization epitope on the gp120 glycoprotein of human immunodeficiency virus type 1. J Virol, 70(2), 1100-1108. <u>PUBMED</u>

NOTE: Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Anti-HIV-1 gp120 Monoclonal (2G12) from Polymun Scientific." Also include the references cited above in any publications.

Last Updated June 13, 2017

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.