



## NIH AIDS Reagent Program

20301 Century Boulevard  
Building 6, Suite 200  
Germantown, MD 20874  
USA

Phone: 240 686 4740  
Fax: 301 515 4015  
aidsreagent.org

### DATA SHEET

**Reagent:** HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 48D TK)

**Catalog Number:** 2344

**Lot Number:** 180423

**Release Category:** B

**Provided:** 5 µg of dried purified DNA stabilized in DNASTable *Plus*

**Cloning Site:** The size of the insert is approximately 144 bp.

**Cloning Vector:** pGEX2TK  
Ampicillin resistant

**Description:** An expression vector which produces GST fused with HIV-1 HXB2 Tat protein that has been truncated after aa 48. The wildtype Tat activation domain is expressed.

**Special Characteristics:** This construct is 5292 bp including the insert.

This plasmid contains a thrombin proteolytic site between the *Schistosoma japonicum* glutathione S-transferase (GST) sequence and the Tat sequence. Tat is expressed as a GST fusion protein and can be cleaved and purified by thrombin proteolytic digestion.

This construct contains the phosphorylation site for the catalytic subunit of camp-dependent heart muscle kinase. GST-Tat fusions in this vector can thus be labeled directly in vitro to high specific activities with (γ<sup>32</sup>P)ATP and commercially available kinase.

[Contributor provided sequence file](#)

[Sequence file lot 180423](#)

Additional HIV-1 and HIV-2 GST-Tat expression vectors are also available. [GST-Tat Expression Vectors](#)

Plasmids can be propagated in STBL2 cells and grown at 37°C. Larger plasmids may benefit from growth at 30°C. This construct may also be grown in other competent

---

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

cells.

This reagent is currently being provided as dried purified DNA stabilized in DNASTable *PLUS*. Please see the notice for additional information and the protocol for reconstitution of dried DNA reagents. [Dried DNA Notice](#)

**Recommended Storage:** Keep the reagent at room temperature in a dry storage cabinet or in a moisture barrier bag.

**Contributor:** Dr. Andrew Rice

**References:**

Herrmann, C. H. and Rice, A. P. (1993). Specific interaction of the human immunodeficiency virus Tat proteins with a cellular protein kinase. *Virology*, 197(2), 601-8. doi:10.1006/viro.1993.1634 [PUBMED](#)

Herrmann, C. H. and Rice, A. P. (1995). Lentivirus Tat proteins specifically associate with a cellular protein kinase, TAK, that hyperphosphorylates the carboxyl-terminal domain of the large subunit of RNA polymerase II: candidate for a Tat cofactor. *J Virol*, 69(3), 1612-20. [PUBMED](#)

Rhim, H., Echetebe, C. O., Herrmann, C. H. and Rice, A. P. (1994). Wild-type and mutant HIV-1 and HIV-2 Tat proteins expressed in *Escherichia coli* as fusions with glutathione S-transferase. *J Acquir Immune Defic Syndr*, 7(11), 1116-21. [PUBMED](#)

**NOTE:** Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 48D TK) from Dr. Andrew Rice (cat# 2344)." Also include the references cited above in any publications.

**Scientists at for-profit institutions or who intend commercial use of this reagent must contact the Baylor College of Medicine at the following email addresses: [mta@bcm.edu](mailto:mta@bcm.edu) and [blg@bcm.edu](mailto:blg@bcm.edu), before the reagent can be released.**

**Last Updated:** March 24, 2020

---

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.