



NIH AIDS Reagent Program

20301 Century Boulevard
Building 6, Suite 200
Germantown, MD 20874
USA

Phone: 240 686 4740
Fax: 301 515 4015
aidsreagent.org

DATA SHEET

Reagent: HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 48D C22G)

Catalog Number: 2361

Lot Number: 190017

Release Category: B

Provided: 5 µg of dried purified DNA stabilized in DNastable PLUS

Cloning Site: The size of the insert is approximately 144 bp.

Cloning Vector: pGEX2T
Ampicillin resistant

Description: An expression vector which produces GST fused with HIV-1 HXB2 Tat protein that has been truncated after aa 48. The non-functional Tat activation domain is expressed.

Special Characteristics: This construct is 5094 bp including the insert.
This plasmid contains a thrombin proteolytic site between the Schistosoma japonicum glutathione S-transferase (GST) sequence and the Tat sequence. Tat is expressed as a GST fusion protein and can be cleaved and purified by thrombin proteolytic digestion.
[Contributor provided sequence file](#)
[Sequence file lot 190017](#)
Additional HIV-1 and HIV-2 GST-Tat expression vectors are also available. [GST-Tat Expression Vectors](#)
Plasmids can be propagated in STBL2 cells and grown at 37°C. Larger plasmids may benefit from growth at 30°C. This construct may also be grown in other competent cells.
This reagent is currently being provided as dried purified DNA stabilized in DNastable PLUS. Please see the notice for additional information and the protocol for reconstitution of dried DNA reagents. [Dried DNA Notice](#)

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

reconstitution of dried DNA reagents. [Dried DNA NOTICE](#)

Recommended Storage:

Keep the reagent at room temperature in a dry storage cabinet or in a moisture barrier bag.

Contributor:

Dr. Andrew Rice

References:

Herrmann, C. H. and Rice, A. P. (1993). Specific interaction of the human immunodeficiency virus Tat proteins with a cellular protein kinase. *Virology*, 197(2), 601-8. doi:10.1006/viro.1993.1634 [PUBMED](#)

Herrmann, C. H. and Rice, A. P. (1995). Lentivirus Tat proteins specifically associate with a cellular protein kinase, TAK, that hyperphosphorylates the carboxyl-terminal domain of the large subunit of RNA polymerase II: candidate for a Tat cofactor. *J Virol*, 69(3), 1612-20. [PUBMED](#)

Rhim, H., Echetebe, C. O., Herrmann, C. H. and Rice, A. P. (1994). Wild-type and mutant HIV-1 and HIV-2 Tat proteins expressed in *Escherichia coli* as fusions with glutathione S-transferase. *J Acquir Immune Defic Syndr*, 7(11), 1116-21. [PUBMED](#)

NOTE:

Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 48D C22G) from Dr. Andrew Rice (cat# 2361)." Also include the references cited above in any publications.

Scientists at for-profit institutions or who intend commercial use of this reagent must contact the Baylor College of Medicine at the following email addresses: mta@bcm.edu and blg@bcm.edu, before the reagent can be released.

Last Updated:

March 24, 2020

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.