



NIH AIDS Reagent Program

20301 Century Boulevard
Building 6, Suite 200
Germantown, MD 20874
USA

Phone: 240 686 4740
Fax: 301 515 4015
aidsreagent.org

DATA SHEET

Reagent: HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 (72R) C22G)

Catalog Number: 2362

Lot Number: 180451

Release Category: B

Provided: 5 µg of dried purified DNA stabilized in DNASTable *Plus*

Cloning Site: The size of the insert is approximately 216 bp.

Cloning Vector: pGEX2T
Ampicillin resistant

Description: An expression vector which produces GST fused with the first exon of transactivation negative HIV-1 HXB2 Tat protein that contains a C22G mutation.

Special Characteristics: This construct is 5163 bp including the insert.
This plasmid contains a thrombin proteolytic site between the Schistosoma japonicum glutathione S-transferase (GST) sequence and the Tat sequence. Tat is expressed as a GST fusion protein and can be cleaved and purified by thrombin proteolytic digestion.
[Contributor provided sequence file](#)
[Sequence file lot 180451](#)
Additional HIV-1 and HIV-2 GST-Tat expression vectors are also available. [GST-Tat Expression Vectors](#)
Plasmids can be propagated in STBL2 cells and grown at 37°C. Larger plasmids may benefit from growth at 30°C. This construct may also be grown in other competent cells.
This reagent is currently being provided as dried purified DNA stabilized in DNASTable *PLUS*. Please see the notice for additional information and the protocol for reconstitution of dried DNA reagents. [Dried DNA Notice](#)

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

reconstitution of dried DNA reagents. [Dried DNA NOTICE](#)

Recommended Storage:

Keep the reagent at room temperature in a dry storage cabinet or in a moisture barrier bag.

Contributor:

Dr. Andrew Rice

References:

Herrmann, C. H. and Rice, A. P. (1993). Specific interaction of the human immunodeficiency virus Tat proteins with a cellular protein kinase. *Virology*, 197(2), 601-8. doi:10.1006/viro.1993.1634 [PUBMED](#)

Herrmann, C. H. and Rice, A. P. (1995). Lentivirus Tat proteins specifically associate with a cellular protein kinase, TAK, that hyperphosphorylates the carboxyl-terminal domain of the large subunit of RNA polymerase II: candidate for a Tat cofactor. *J Virol*, 69(3), 1612-20. [PUBMED](#)

Rhim, H., Echetebe, C. O., Herrmann, C. H. and Rice, A. P. (1994). Wild-type and mutant HIV-1 and HIV-2 Tat proteins expressed in *Escherichia coli* as fusions with glutathione S-transferase. *J Acquir Immune Defic Syndr*, 7(11), 1116-21. [PUBMED](#)

NOTE:

Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 HXB2 GST-Tat Expression Vector (GST-Tat 1 (72R) C22G) from Dr. Andrew Rice (cat# 2362)." Also include the references cited above in any publications.

Scientists at for-profit institutions or who intend commercial use of this reagent must contact the Baylor College of Medicine at the following email addresses: mta@bcm.edu and blg@bcm.edu, before the reagent can be released.

Last Updated:

March 26, 2020

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.