



NIH AIDS Reagent Program

20301 Century Boulevard
Building 6, Suite 200
Germantown, MD 20874
USA

Phone: 240 686 4740
Fax: 301 515 4015
aidsreagent.org

DATA SHEET

Reagent:	☒ HIV-1 RF/41-D2 Virus
Catalog Number:	2805
Lot Number:	099076
Release Category:	A
Provided:	1 mL cell-free virus TCID ₅₀ = 2.2 x 10 ⁴ /mL
Host Strain:	MT-2
Propagation:	Propagated in MT-2 cells.
Sterility:	Negative for mycoplasma, bacteria and fungi.
Description:	Protease mutant derived from HIV-1 RF strain.
Special Characteristics:	Single and double protease gene mutant strains were selected <i>in vitro</i> in the presence of the novel protease inhibitors DMP323 (a cyclic urea protease inhibitor) or P9941 (a C-2 symmetrical diol protease inhibitor). Four mutants derived from HIV-1 _{RF} and one from the AZT-resistant strain Donor E are available. All of those viruses are stable in culture, and do not require the addition of antivirals to maintain phenotype. All are extremely cytopathic and require frequent feeding with uninfected cells. Table of additional HIV-1 Protease mutants included in this set.
Recommended Storage:	Liquid nitrogen.
Contributor:	Dr. Dean Winslow.

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

References: Otto, M. J., S. Garber, D. L. Winslow, C. D. Reid, P. Aldrich, P. K. Jadhav, C. E. Patterson, C. N. Hodge and Y. S. Cheng. (1993). In vitro isolation and identification of human immunodeficiency virus (HIV) variants with reduced sensitivity to C-2 symmetrical inhibitors of HIV type 1 protease. *Proc Natl Acad Sci U S A*, 90(16), 7543-7. [PUBMED](#)

King RW, Garber S, Winslow DL, Reid C, Bacheler LT, Anton E, Otto MJ. Multiple mutations in the human immunodeficiency virus protease gene are responsible for decreased susceptibility to protease inhibitors. *Antiviral Chem Chemother* **6**:80-88, 1995.

Otto MJ, Reid CD, King RW, Garber S, Baker DB, Anton E, Winslow DL. Exposure of chronically infected PBMC's to DMP450 can completely suppress virus replication or select resistant variants depending upon the dose of compound. *Second National Conference on Human Retroviruses and Related Infections*, Washington, DC, January 29-February 2, 1995.

NOTE: Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 RF/41-D2 Virus from Dr. Dean Winslow." Also include the references cited above in any publications.

Last Updated: November 14, 2018

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.