



NIH AIDS Reagent Program

20301 Century Boulevard
Building 6, Suite 200
Germantown, MD 20874
USA

Phone: 240 686 4740
Fax: 301 515 4015
aidsreagent.org

DATA SHEET

Reagent:	Anti-HIV-1 p24 Gag Monoclonal (#24-2)
Catalog Number:	6457
Lot Number:	060883
Release Category:	A
Provided:	286 μ l (1.2 mg) affinity purified antibody at 4.2 mg/ml.
Host:	Balb/c splenocyte x SP2 myeloma.
Special Characteristics:	<p>Affinity purified from ascites fluid by Protein A/G chromatography. The storage buffer is phosphate-buffered saline (0.02% sodium azide). ELISA titer against HIV-1IIIB p24 lysate (50ng/well) is 1:100,000.</p> <p><i>Immunogen:</i> Hexahistidine, amino-terminal tagged HIV-1 (HXB-3 isolate) p24 Gag protein, expressed in <i>E. coli</i>, purified under denaturing conditions by chelate-affinity chromatography and dialyzed into a phosphate/salt buffer (pH 10).</p> <p><i>Immunization:</i> 5 to 50 μg of immunogen was injected intraperitoneally per mouse as an emulsion with complete Freund's adjuvant. Subsequent booster immunizations (at least three) using incomplete Freund's adjuvant were also intraperitoneal and were administered every three weeks starting at week two. A final boost of immunogen in phosphate buffered saline was administered intravenously (tail vein) three days prior to removal of the spleen.</p> <p><i>Applications:</i> This antibody has been used successfully for Western blots (HXB-3 and YU-2 isolates) and immunofluorescence assays. Its epitope has not been mapped.</p>
Recommended Storage:	-70°C.
Contributor:	Dr. Michael H. Malim.
Isotype:	IgG ₁ , κ .

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

References:

Simon JH, Fouchier RA, Southerling TE, Guerra CB, Grant CK, Malim MH. The Vif and Gag proteins of human immunodeficiency virus type 1 colocalize in infected human T cells. *J Virol* **71**:5259-5267, 1997.

Fouchier RA, Meyer BE, Simon JH, Fischer U, Malim MH. HIV-1 infection of non-dividing cells: evidence that the amino-terminal basic region of the viral matrix protein is important for Gag processing but not for post-entry nuclear import. *EMBO J* **16**:4531-4539, 1997.

NOTE:

Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 p24 Gag Monoclonal (#24-2) from Dr. Michael H. Malim." Also include the references cited above in any publications.

Last Updated

February 02, 2018

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.