

## NIH AIDS Reagent Program

20301 Century Boulevard Building 6, Suite 200 Germantown, MD 20874 USA Phone: 240 686 4740 Fax: 301 515 4015 aidsreagent.org

## **DATA SHEET**

**Reagent:** Anti-HIV-1 gp160 Hybridoma (Chessie 13-39.1)

Catalog Number: 990

Lot Number: 011252

Release Category: B

**Propagation** 

**Provided:**  $4 \times 10^6$  cells/ml. Viability is 96%.

**Medium:** μg/ml streptomycin, 90%; fetal bovine serum, 10%.

**Freeze Medium:** Fetal bovine serum, 90%; DMSO, 10%.

**Growth**Thaw cells quickly in a 37°C water bath and dilute with 15 ml of propagation medium warmed to 37°C. Centrifuge to remove DMSO. For exponential growth, maintain the

warmed to 37°C. Centrifuge to remove DMSO. For exponential growth, maintain the cells at  $2-5 \times 10^5/\text{ml}$  and split every 1-2 days. Doubling time is 14 hours. Cells are hardy. Freeze cells at  $10^6-10^7/\text{ml}$ . To obtain antibody, overgrow cells until they die.

RMPI 1640 supplemented with 50 µg/ml \(\beta\)-mercaptoethanol, 100 U/ml penicillin, 100

Yields about 10-20 µg antibody/ml.

**Description** Balb/c mouse splenocyte x P3X63 Ag8.X653 myeloma

**Special** Chessie 13-39.1 produces an IgG1 monoclonal antibody that maps to aa 252-273 **Characteristics:** (RPVVSTQLLLNGSLAEEEVVIR) of LAI qp120. The antibody recognizes epitopes wh

(RPVVSTQLLLNGSLAEEEVVIR) of LAI gp120. The antibody recognizes epitopes which survive alkylation and reduction. Antibody reactivities have been tested using the

HIV-III<sub>R</sub> isolate and its cloned env gene products.

**Sterility:** Negative for mycoplasma, bacteria and fungi.

Recommended

Storage:

Liquid nitrogen.

Contributor: Dr. George K. Lewis.

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

REV: 06/05/2017 Page 1 of 2

References: Abacioglu YH, Fouts TR, Laman JD, Claassen E, Pincus SH, Moore JP, Roby CA,

Kamin-Lewis R, Lewis GK. Epitope mapping and topology of baculovirus-expressed HIV-1 gp160 determined with a panel of murine monoclonal antibodies. AIDS Res Hum

Retroviruses 10:371-381, 1994.

NOTE:

Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Anti-HIV-1 gp160 Hybridoma (Chessie 13-39.1) from Dr. George Lewis." Also include the reference

cited above in any publications.

June 05, 2017 **Last Updated** 

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

REV: 06/05/2017 Page 2 of 2