

Human Metapneumovirus, TN/83-1211

Catalog No. NR-22227

Product Description:

Human metapneumovirus (HMPV), TN/83-1211 was isolated in 1983 from a human specimen collected in Tennessee, USA. NR-22227 lot 70039699 was produced by infecting *Macaca mulatta* kidney epithelial cells (LLC-MK2; ATCC® CCL-7.1™) in Opti-MEM® Minimal Essential Medium (Gibco® 31985) supplemented with 2 mM L-glutamine (ATCC® 30-2214™), 100 µg per mL CaCl₂ and 5 µg per mL trypsin (ATCC® 30-2101™) for 5 days at 37°C and 5% CO₂.

Passage History:

L7/L2 (Vanderbilt/BEI Resources); L = LLC-MK2 cells

Lot: 70039699

Manufacturing Date: 30NOV2020

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in LLC-MK2 Derivative Cells	Report results	Rounding, sloughing and syncytia formation
Identification by Direct Fluorescent Antibody (DFA) Assay ¹	Fluorescence observed	Fluorescence observed
Sequencing of Species-Specific Region (~ 880 nucleotides)	≥ 98% identity with HMPV, TN/83-1211 complete genome (GenBank: KC562244.1)	99.7% identity with HMPV, TN/83-1211 complete genome (GenBank: KC562244.1) ²
Titer by TCID ₅₀ Assay in LLC-MK2 Cells with DFA Readout ^{1,3} (7 days at 37°C and 5% CO ₂)	Report results	1.58 × 10 ⁶ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ⁴ Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Using Light Diagnostics™ Human Metapneumovirus DFA Reagent (Millipore 5091RUO)

²The limited nucleotide sequencing of NR-22227 performed at BEI Resources is not sufficient to confirm exact strain identity owing to the high degree of sequence conservation within HMPV lineages.

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

17 MAR 2021

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

