

Product Information Sheet for NR-210

***Yersinia enterocolitica* subsp. *enterocolitica*, Strain WA-314**

Catalog No. NR-210

(Derived from ATCC® 51871™)

For research only. Not for use in humans.

Contributor:

ATCC®

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Enterobacteriaceae*, *Yersinia*

Species: *Yersinia enterocolitica* subsp. *enterocolitica*^{1,2}

Strain: WA-314

Biotype: 1B

Serotype: O:8

Original Source: *Yersinia enterocolitica* (*Y. enterocolitica*) subsp. *enterocolitica*, strain WA-314 is a human clinical specimen.³

Comments: *Y. enterocolitica* subsp. *enterocolitica*, strain WA-314 was deposited at ATCC® by Dr. Alexander V. Rakin, Max von Pettenkofer-Institute of Hygiene and Medical Microbiology, University of Munich, Munich, Germany. The complete genome of *Y. enterocolitica* subsp. *enterocolitica*, strain WA-314 is available (GenBank: [NZ_AKKR000000000.1](https://www.ncbi.nlm.nih.gov/nuclseq/NC_009411.1)).

Y. enterocolitica subsp. *enterocolitica* is a significant food-borne enteropathogen which causes gastroenteritis.⁴ *Y. enterocolitica* subsp. *enterocolitica* is an extremely heterogeneous species, encompassing six biotypes and currently more than 50 serogroups, not all of which can cause disease.⁵ It is of particular concern to the food industry because it is a psychrotrophic pathogen able to proliferate at temperatures approaching 0°C.

Y. enterocolitica subsp. *enterocolitica* is a non-spore-forming, Gram-negative, rod-shaped coccobacillus. Virulence-associated genes are located on the chromosome and on the pYV (approximately 70 kb) plasmid found in typical virulent strains of *Y. enterocolitica* subsp. *enterocolitica*.^{6,7} This plasmid encodes a type III secretion system involved in the delivery of virulence proteins that contribute to internalization into host cells.⁷

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Tryptic Soy Broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-210 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored

at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Brain Heart Infusion broth or Tryptic Soy broth or equivalent

Tryptic Soy agar or Sheep Blood agar or equivalent

Incubation:

Temperature: 26°C⁸

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; thaw slowly.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 26°C for 1 to 2 days.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Yersinia enterocolitica* subsp. *enterocolitica*, Strain WA-314, NR-210."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 6th ed. Washington, DC: U.S. Government Printing Office, 2020; see www.cdc.gov/biosafety/publications/bmbl5/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for

damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. "Validation List No. 75." *Int. J. Syst. Evol. Microbiol.* 50 (2000): 1415-1417.
2. Neubauer, H., et al. "Yersinia enterocolitica 16S rRNA Gene Types Belong to the Same Genospecies but Form Three Homology Groups." *Int. J. Med. Microbiol.* 290 (2000): 61-64. PubMed: 11043982.
3. Heesemann, J. "Chromosomal-Encoded Siderophores Are Required for Mouse Virulence of Enteropathogenic Yersinia Species." *FEMS Microbiol. Lett.* 48 (1987): 229-233.
4. Fredriksson-Ahomaa, M., A. Stolle and H. Korkeala. "Molecular Epidemiology of Yersinia enterocolitica Infections." *FEMS Immunol. Med. Microbiol.* 47 (2006): 315-329. PubMed: 16872368.
5. Viridi, J. S. and P. Sachdeva. "Molecular Heterogeneity in Yersinia enterocolitica and 'Y. enterocolitica-Like' Species – Implications for Epidemiology, Typing and Taxonomy." *FEMS Immunol. Med. Microbiol.* 45 (2005): 1-10. PubMed: 15985218.
6. Bottone, E. J. "Yersinia enterocolitica: The Charisma Continues." *Clin. Microbiol. Rev.* 10 (1997): 257-276. PubMed: 9105754.
7. Snellings, N. J., M. Popek and L. E. Lindler. "Complete DNA Sequence of Yersinia enterocolitica Serotype O:8 Low-Calcium-Response Plasmid Reveals a New Virulence Plasmid-Associated Replicon." *Infect. Immun.* 69 (2001): 4627-4638. PubMed: 11402007.
8. Chu, M. C. *Laboratory Manual of Plague Diagnostic Tests*. Centers for Disease Control and Prevention, Atlanta, 2000.
9. Iwobi, A., et al. "Novel Virulence-Associated Type II Secretion System Unique to High-Pathogenicity Yersinia enterocolitica." *Infect. Immun.* 71 (2003): 1872-1879. PubMed: 12654803.
10. Iwobi, A., et al. "Representational Difference Analysis Uncovers a Novel IS10-Like Insertion Element Unique to Pathogenic Strains of Yersinia enterocolitica." *FEMS Microbiol. Lett.* 210 (2002): 251-255. PubMed: 12044683.
11. Trülsch, K., et al. "Cloning and Characterization of the Gene Encoding Periplasmic 2',3'-Cyclic Phosphodiesterase of Yersinia enterocolitica O:8." *Microbiology* 147 (2001): 203-213. PubMed: 11160814.
12. Rakin, A., et al. "Local Hopping of IS3 Elements into the A+T-Rich Part of the High-Pathogenicity Island in Yersinia

enterocolitica 1B, O:8." *FEMS Microbiol. Lett.* 182 (2000): 225-229. PubMed: 10620670.

13. Rakin, A., et al. "Common and Specific Characteristics of the High-Pathogenicity Island of Yersinia enterocolitica." *Infect. Immun.* 67 (1999): 5265-5274. PubMed: 10496905.
14. Rakin, A., et al. "The Pesticin Receptor of Yersinia enterocolitica: A Novel Virulence Factor with Dual Function." *Mol. Microbiol.* 13 (1994): 253-263. PubMed: 7984105.
15. Heesemann, J., et al. "Virulence of Yersinia enterocolitica is Closely Associated with Siderophore Production, Expression of an Iron-Repressible Outer Membrane Polypeptide of 65,000 Da and Pesticin Sensitivity." *Mol. Microbiol.* 8 (1993): 397-408. PubMed: 8316088.

ATCC® is a trademark of the American Type Culture Collection.

