

Certificate of Analysis for NR-9696

Influenza B Virus, B/Florida/4/2006 (Yamagata Lineage)

Catalog No. NR-9696

This reagent is the tangible property of the U.S. Government.

Product Description:

Influenza B virus, B/Florida/4/2006 [Yamagata Lineage (BY)] from a human in Florida, USA on November 1, 2006. NR-9696 lot 70051034 was produced in the allantoic cavity of specific pathogen-free (SPF) embryonated chicken eggs (9-to 11-day-old; Charles River, Norwich, Connecticut, USA) infected with seed material lot 58155279 for 3 days at 33°C in a humidified chamber.

Lot: 70051034 Manufacturing Date: 17MAR2022

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin Coding Regions (~ 900 nucleotides)	≥ 98% identity with B/Florida/4/2006 (Yamagata Lineage) (GenBank: CY073895)	100% identity with B/Florida/4/2006 (Yamagata Lineage) (GenBank: CY073895)
Titer by CEID ₅₀ Assay in Embryonated Chicken Eggs ¹ (3 days at 33°C in a humidified chamber)	Report results	1.6 × 10 ⁸ CEID ₅₀ per mL
Sterility (21-day incubation)		
Harpo's HTYE broth, 37°C and 26°C, aerobic ²	No growth	No growth
Trypticase Soy broth, 37°C and 26°C, aerobic	No growth	No growth
Sabouraud broth, 37°C and 26°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, anaerobic	No growth	No growth
Thioglycollate broth, 37°C, anaerobic	No growth	No growth
DMEM with 10% FBS, 37°C, aerobic	No growth	No growth
Mycoplasma Contamination		
Agar and broth culture (14-day incubation at 37°C)	None detected	None detected
DNA detection by PCR of extracted Test Article nucleic acid	None detected	None detected

¹The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

/Sonia Bjorum Brower/ Sonia Bjorum Brower

25 MAY 2022

Lead Technical Writer, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org Tel: 800-359-7370

Fax: 703-365-2898

²Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.