

**Human Coxsackievirus B1, Conn-5**

**Catalog No. NR-58650**

This reagent is the property of the U.S Government

**Product Description:**

Human coxsackievirus B1 (hCVB1), Conn-5 was isolated from the stool of a patient with aseptic meningitis in Connecticut, USA, in 1948. NR-58650 lot 70054069 was produced by infecting *Macaca mulatta* (Rhesus monkey) kidney epithelial cells (LLC-MK2 derivative; ATCC® CCL-7.1™) with the deposited material and incubating in Eagle's Minimum Essential Medium (ATCC® 30-2003™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 3 days at 37°C with 5% CO<sub>2</sub>.

**Passage History:**

RhMK(7)AGMK(1)/AGMK(1)RhMK(1) (Prior to deposit/BEI Resources); RhMK = *Macaca mulatta* (Rhesus monkey) kidney epithelial cells; AGMK = *Cercopithecus aethiops* (*African green*) monkey kidney epithelial cells

**Lot: 70054069**

**Manufacturing Date: 12AUG2022**

TEST	SPECIFICATIONS	RESULTS
<b>Identification by Infectivity in LLC-MK2 Cells</b>	Cell rounding and detachment	Cell rounding and detachment
<b>Sequencing of Species-Specific Region</b> Polyprotein 2 (~ 640 nucleotides)	≥ 98% identity with hCVB1, complete genome (GenBank: M16560.1)	100% identity with hCVB1, complete genome (GenBank: M16560.1)
<b>Titer by TCID<sub>50</sub> Assay in LLC-MK2 Cells by Cytopathic Effect<sup>1</sup></b> (9 days at 37°C with 5% CO <sub>2</sub> )	Report results	5.0 × 10 <sup>8</sup> TCID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>2</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>2</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorum Brower/

Sonia Bjorum Brower

26 JAN 2022

Lead Technical Writer or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

