

**Genomic RNA from SARS-Related Coronavirus 2, Isolate hCoV-19/USA/COR-22-063113/2022 (Lineage BA.5; Omicron Variant)**

**Catalog No. NR-58496**

**Product Description:**

Genomic RNA was isolated from a preparation of cell lysate and supernatant from *Homo sapiens* lung adenocarcinoma epithelial cells (Calu-3; ATCC® HTB-55™) infected with SARS-CoV-2, isolate hCoV-19/USA/COR-22-063113/2022. (BEI Resources lot 70053501) using QIAamp® Viral RNA Mini Kit (Qiagen® 52906). The viral genomic RNA is in a background of cellular nucleic acid and carrier RNA.

**Lot: 70056168**

**Manufacturing Date: 29SEP2022**

TEST	SPECIFICATIONS	RESULTS
<b>Genotypic Analysis</b> Sequencing of species-specific region (~ 920 nucleotides)	≥ 98% identity with SARS-CoV-2, hCoV-19/ USA/COR-22-063113/2022 (depositor sequence)	100% identity with SARS-CoV-2, hCoV-19/ USA/COR-22-063113/2022 (depositor sequence)
<b>Functional Activity by RT-PCR Amplification<sup>1</sup></b> ORF1ab gene	~ 950 base pair amplicon	~ 950 base pair amplicon (Figure 1)
<b>Estimated Concentration (post-dilution) by Qubit® Measurement (Viral, Cellular and Carrier)<sup>2</sup></b>	Report results	0.6 ng per 100 µL (0.006 µg/mL)
<b>Estimated Amount per Vial<sup>2</sup></b>	Report results	0.6 ng
<b>NAAT Detectable Units Using BioRad QX200 Droplet Digital PCR (ddPCR™) System<sup>3</sup></b> (Post vial; 8 replicates)	Report results	2.85× 10 <sup>6</sup> NDU per mL
<b>Virus Inactivation</b> 10% of total yield inoculated on Calu-3 cells and evaluated for cytopathic effect and by qPCR after serial passage <sup>4</sup>	No viable virus detected	No viable virus detected


<sup>1</sup>Amplified using iTaq™ Universal SYBR Green One-step Kit (Bio-Rad® 172-5151) with 5 µL of NR-58496 in a 50 µL reaction

<sup>2</sup>Measurement is determined pre-vial prior to dilution due to the limit of detection of the quantification method

<sup>3</sup>NDU; NAAT(Nucleic acid amplification testing) Detectable Unit

<sup>4</sup>Use of the QIAamp® Viral RNA Mini Kit has been demonstrated to consistently inactivate 100% of SARS-CoV-2 as shown by the absence of cytopathic effect (CPE) and viral RNA expression by qPCR after plating the entire extract on virus-susceptible cells for two passages.

**Figure 1: Functional Activity of NR-58496 by RT-PCR Amplification of ORF1ab Gene**



Lane 1: Invitrogen™ TrackIt™ 1 Kb Plus DNA Ladder  
Lane 2: PCR product from 1 µL of NR-58496

/Sonia Bjorum Brower/

Sonia Bjorum Brower

28 FEB 2023

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

