

Influenza B Virus, B/New York/1055/2003

Catalog No. NR-48660

Product Description:

Influenza B virus, B/New York/1055/2003 was isolated on January 29, 2003, from an adult male human in Tompkins County, New York, USA. NR-48660 lot 70059265 was produced by infecting Madin-Darby Canine Kidney cells (MDCK; ATCC® CCL-34™) with influenza B virus, B/New York/1055/2003 and incubating in Eagle's Minimum Essential Medium (ATCC® 30-2003™) supplemented with 0.125% bovine serum albumin and 1 µg/mL L-1-tosylamido-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin for 5 days at 33°C and 5% CO₂.

Passage History:

X(?) / C(4) (Prior to deposit at BEI Resources/BEI Resources); C = MDCK cells

Lot: 70059265

Manufacturing Date: 15MAR2023

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in MDCK Cells	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Neuraminidase Coding Regions (~ 910 nucleotides)	≥ 98% identity with B/New York/1055/2003 (GenBank: CY174331.1)	100% identity with B/New York/1055/2003 (GenBank: CY174331.1)
Titer by TCID₅₀ Assay in MDCK Cells by CPE¹ (8 days at 37°C and 5% CO ₂)	Report results	8.9 × 10 ⁶ TCID ₅₀ /mL
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ² Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic	None detected None detected	None detected None detected

¹The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

²Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorum Brower/

Sonia Bjorum Brower

17 AUG 2023

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

