SUPPORTING INFECTIOUS DISEASE RESEARCH

# Influenza A Virus, A/Sydney/5/1997 (H3N2)

### Catalog No. NR-12278

This reagent is the property of the U.S. Government.

### **Product Description:**

Influenza A virus, A/Sydney/5/1997 (H3N2) was isolated from human in Sydney, Australia in 1997. NR-12278 lot 70060721 was produced in the allantoic cavity of specific pathogen free (SPF) embryonated chicken eggs (10- to 11- day-old; Charles River, Norwich, Connecticut, USA) infected with the deposited material for 2 days at 35°C in a humidified chamber.

## Lot: 70060721

# Manufacturing Date: 01JUN2023

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells		Positive
Sequencing of Hemagglutinin and Matrix Coding Regions Matrix (~ 940 nucleotides)	≥ 98% identity with A/Sydney/5/1997 (H3N2) (GenBank CY039080)	99.8% identity with A/Sydney/5/1997 (H3N2) (GenBank CY039080)
Titer by CEID₅₀ Assay in Embryonated Chicken Eggs¹ (2 days at 35°C in a humidified chamber)	Report results	5 × 10 <sup>6</sup> CEID <sub>50</sub> /mL
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>2</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>The Chicken Embryo Infectious Dose 50% (CEID<sub>50</sub>) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID<sub>50</sub> provides a measure of the infectious titer (or infectivity) of a virus preparation.

<sup>2</sup>Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

# /Sonia Bjorum Brower/

Sonia Bjorum Brower

Technical Manager or designee, ATCC Federal Solutions

ATCC<sup>®</sup>, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC<sup>®</sup>'s knowledge.



05 OCT 2023

ATCC<sup>®</sup> is a trademark of the American Type Culture Collection. You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org