

**Influenza A Virus, A/Baltimore/JH-335/2022 (H3N2)**

**Catalog No. NR-59460**

**Product Description:**

Influenza A virus, A/Baltimore/JH-335/2022 (H3N2) was isolated from a human in Baltimore, Maryland, USA, in 2022. NR-59460 lot 70062479 was produced by infecting Madin-Darby canine kidney-SIAT1 cells (MDCK-SIAT1; BEI Resources lot 70047595) with influenza A virus, A/Baltimore/JH-335/2022 (H3N2) and incubating in Dulbecco's Modified Eagle Medium (ATCC® 30-2002™) supplemented with 0.3% bovine serum albumin and 5 µg/mL N-acetyl trypsin for 4 days at 33°C and 5% CO<sub>2</sub>. The cells and supernatant were spin-clarified at 500 × g for 10 minutes at 4°C.

**Passage History:**

hNEC(1), MDCK-SIAT1(1) / MDCK-SIAT1(1) (Johns Hopkins University/BEI Resources); hNEC = human Nasal Epithelial Cells; MDCK-SIAT = Madin-Darby canine kidney-SIAT1 cells

**Lot: 70062479**

**Manufacturing Date: 10SEP2023**

TEST	SPECIFICATIONS	RESULTS
<b>Identification by Infectivity in MDCK-SIAT1 Cells</b>	Cell rounding and detachment	Cell rounding and detachment
<b>Sequencing of Hemagglutinin and Neuraminidase Coding Regions</b> Hemagglutinin (~ 1740 nucleotides) Neuraminidase (~ 1440 nucleotides)	Consistent with hemagglutinin type 3 (H3) Consistent with neuraminidase type 2 (N2)	Consistent with H3 Consistent with N2
<b>Next-Generation Sequencing (NGS) of Complete Genome Using Illumina® MiSeq™ Platform</b>	Consistent with sequence of depositor's material	Consistent with sequence of depositor's material
<b>Titer by TCID<sub>50</sub> Assay in MDCK-SIAT1 Cells by Hemagglutination Assay<sup>1,2</sup></b> (7 days at 33°C and 5% CO <sub>2</sub> )	Report results	8.9 × 10 <sup>7</sup> TCID <sub>50</sub> /mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>3</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the infectious titer (or infectivity) of a virus preparation.

<sup>2</sup>Assay performed using 0.5% turkey red blood cells

<sup>3</sup>Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorum Brower/  
Sonia Bjorum Brower

Technical Manager or designee, ATCC Federal Solutions

16 JAN 2024

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

