

Streptococcus pneumoniae, Strain OREP17F

Catalog No. NR-59136

For research use only. Not for use in humans.

Contributor:

Moon H. Nahm, M.D., Professor, Department of Pathology, University of Alabama at Birmingham, Alabama, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: Streptococcaceae, Streptococcus

Species: Streptococcus pneumoniae

Strain: OREP17F

Source: Streptococcus pneumoniae (S. pneumoniae), strain OREP17F was derived from a human wild-type S. pneumoniae, strain DS3022-06 (serotype 17F) by natural selection using increasing concentrations of optochin.^{1,2}

Comments: S. pneumoniae, strain OREP17F is reported to be resistant to optochin at 2 µg/mL.²

S. pneumoniae is a Gram-positive, α-hemolytic diplococcal aerotolerant anaerobe that is a major cause of pneumonia, bacterial meningitis and otitis media. S. pneumoniae has a polysaccharide capsule that acts as a virulence factor for the organism. There are over ninety different capsular types of S. pneumoniae which differ in virulence, prevalence, and extent of drug resistance.^{3,4}

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Todd-Hewitt broth containing 0.5% (w/v) yeast extract and 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-59136 was packaged aseptically in cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy broth or Todd-Hewitt broth containing 0.5% (w/v) yeast extract or equivalent

Tryptic Soy agar or Tryptic Soy agar with 5% defibrinated sheep blood or Todd-Hewitt agar containing 0.5% (w/v) yeast extract or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic with 5% CO₂

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate at 37°C for 1 to 2 days.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Streptococcus pneumoniae, Strain OREP17F, NR-59136."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. [Biosafety in Microbiological and Biomedical Laboratories \(BMBL\)](#). 6th ed. Washington, DC: U.S. Government Printing Office, 2020.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Burton, R. L., Personal Communication.
2. Burton, R. L. and M. H. Nahm. "Development of a Fourfold Multiplexed Opsonophagocytosis Assay for Pneumococcal Antibodies against Additional Serotypes and Discovery of Serological Subtypes in *Streptococcus pneumoniae* Serotype 20." Clin. Vaccine Immunol. 19 (2012): 835-841. PubMed: 22518015.
3. Jedrzejewski, M. J. "Pneumococcal Virulence Factors: Structure and Function." Microbiol. Mol. Biol. Rev. 65 (2001): 187-207. PubMed: 11381099.
4. Habib, M., B. D. Porter and C. Satzke. "Capsular Serotyping of *Streptococcus pneumoniae* Using the Quellung Reaction." J. Vis. Exp. (2014): e51208. PubMed: 24637727.
5. Cywes-Bentley, C., et al. "Antibody to a Conserved Antigenic Target is Protective Against Diverse Prokaryotic and Eukaryotic Pathogens." Proc. Natl. Acad. Sci. USA 110 (2013): E2209-E2218. PubMed: 23716675.
6. Balloch, A., et al. "Interlaboratory Comparison of the Pneumococcal Multiplex Opsonophagocytic Assays and their Level of Agreement for Determination of Antibody Function in Pediatric Sera." mSphere 3 (2018): e00070-18. PubMed: 29695620.
7. Burton, R. L. and M. H. Nahm. "Development and Validation of a Fourfold Multiplexed Opsonization Assay (MOPA4) for Pneumococcal Antibodies." Clin. Vaccine Immunol. 13 (2006): 1004-1009. PubMed: 16960111.

ATCC® is a trademark of the American Type Culture Collection.

