

Clostridioides difficile, Isolate 20110869

Catalog No. NR-49292

Product Description:

Clostridioides difficile (*C. difficile*), isolate 20110869 was obtained from the stool of a young adult male patient with a community-associated (CA) *C. difficile* infection in Tennessee, USA, in 2011. Isolate 20110869 was deposited as PCR ribotype 001_072, North American pulsed-field gel electrophoresis type 2 (NAP2), containing *tcdA*, *tcdB* and *tcdC* of the PaLoc operon. This isolate is reported to be negative for the *C. difficile* binary toxin (CDT). NR-49292 was produced by inoculation of BEI Resources seed lot 63719808 into Modified Reinforced Clostridial broth and incubated for 1 day at 37°C in an anaerobic atmosphere (< 5% O₂; Remel™ Pack-Anaero™). The material from the initial growth was passaged once in Modified Reinforced Clostridial agar kolles for 2 days at 37°C in an anaerobic atmosphere to produce this lot.

Lot: 70065135

Manufacturing Date: 05JAN2024

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology 2 days at 37°C in an anaerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood Motility (wet mount) Hemolysis 2 days at 37°C in an anaerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood VITEK® MS (MALDI-TOF) Biochemical Characterization Esculin hydrolysis ¹ Gelatin hydrolysis ¹	Gram-positive rods Report results Report results Report results <i>C. difficile</i> Positive Positive	Gram-positive rods Irregular, flat, undulate, opaque and gray Non-motile Non-hemolytic <i>C. difficile</i> (99.9%) Positive Positive
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1400 base pairs)	≥ 99% sequence identity to <i>C. difficile</i> type strain (GenBank: MUJVO1000001.1)	100% sequence identity to <i>C. difficile</i> type strain (GenBank: MUJVO1000001.1)
PCR Assay of Extracted DNA Presence of <i>C. difficile</i> -specific genes ² Triose phosphate isomerase (<i>tpi</i>) Presence of toxin genes ^{2,3} <i>cdtB</i> <i>tcdA</i> (wild type) <i>tcdA</i> (partial deletion) <i>tcdB</i>	~ 230 base pair amplicon No amplicon ~ 370 base pair amplicon No amplicon ~ 160 base pair amplicon	~ 230 base pair amplicon No amplicon ~ 370 base pair amplicon No amplicon ~ 160 base pair amplicon
Purity (post-freeze) Anaerobic 7 days at 37°C on Tryptic Soy agar with 5% defibrinated sheep blood Aerobic with 5% CO ₂ 7 days at 37°C on Tryptic Soy agar with 5% defibrinated sheep blood	Report results Report results	Growth consistent with expected colony morphology No growth
Viability (post-freeze) 2 days at 37°C in an anaerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood	Growth	Growth

¹Tests were assessed after 7 days at 37°C in an anaerobic atmosphere.

²Lemee, L., et al. "Multiplex PCR Targeting *tpi* (Triose Phosphate Isomerase), *tcdA* (Toxin A), and *tcdB* (Toxin B) Genes for Toxigenic Culture of *Clostridium difficile*." *J. Clin. Microbiol.* 42 (2004): 5710-5714. PubMed: 15583303.

³Antikainen, J., et al. "Detection of Virulence Genes of *Clostridium difficile* by Multiplex PCR." *APMIS*. 117 (2009): 607-613. PubMed: 19664132.

/Sonia Bjorum Brower/

Sonia Bjorum Brower

26 MAR 2024

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

