

Measles Virus, AIK-C Vaccine Strain

Catalog No. NR-60836

This reagent is the property of the U.S. Government.

Product Description:

Measles virus (MeV), AIK-C vaccine strain was developed from the Edmonston-Enders strain and widely used in Japan. NR-60836 was produced by infecting *Chlorocebus aethiops* kidney epithelial cells with human signaling lymphocytic activation molecule (Vero-hSLAM; BEI Resources NR-55500) with the deposited material and incubating in Dulbecco's Modified Eagle's Medium (ATCC® 30-2002™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 6 days at 37°C with 5% CO₂ to produce this lot.

Passage History:

HK(24)HA(28)/HA(12)SK(33°C)(17)CEF(33°C)(22)¹/Vh(2)/Vh(3) (Initial attenuation to Edmonston-Enders/Prior to deposit at CDC/CDC/BEI Resources); HK = human kidney; HA = human amnion; SK = sheep kidney; CEF = chick embryo fibroblast; Vh= Vero-hSLAM (*Chlorocebus aethiops* kidney epithelial cells with human signaling lymphocytic activation molecule)

Lot: 70079294

Manufacturing Date: 18FEB2026

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero-hSLAM Cells	Cell rounding and detachment	Cell rounding and detachment
Next-Generation Sequencing (NGS) of Complete Genome	≥ 98% sequence identity with MeV, AIK-C vaccine strain (GenBank: AF266286)	99.59% sequence identity with MeV, AIK-C vaccine strain (GenBank: AF266286)
Titer by TCID ₅₀ Assay in Vero-hSLAM Cells by Cytopathic Effect ² (9 days at 37°C with 5% CO ₂)	Report results	2.0 × 10 ³ TCID ₅₀ /mL ³
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ⁴ Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Parks, C. L., et al. "Comparison of Predicted Amino Acid Sequences of Measles Virus Strains in the Edmonston Vaccine Lineage." *J. Virol.* (2001): 910-920. PubMed: 11134304.

²The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

³Titer was determined by cytopathic effects (CPE) and completed in triplicate (2.8 × 10³ TCID₅₀/mL, 2.8 × 10³ TCID₅₀/mL and 2.8 × 10³ TCID₅₀/mL). The average of the three values is reported.

⁴Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorun Brower/

Sonia Bjorun Brower

19 MAY 2026

Technical Manager or designee, ATCC Federal Solutions

Certificate of Analysis for NR-60836

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

