

Influenza A Virus, A/blue-winged teal/Illinois/10OS1546/2010 (H3N6)

Catalog No. NR-35981

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs¹ infected with influenza A virus, A/blue-winged teal/Illinois/10OS1546/2010 (H3N6)

Lot²: 60979639

Manufacturing Date: 14JUN2012

| TEST | SPECIFICATIONS | RESULTS |
|--|--|--|
| Identification by Infectivity Using Embryonated Chicken Eggs¹ Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells | Positive | Positive |
| Sequencing of Hemagglutinin and Matrix Coding Regions Hemagglutinin (608 nucleotides) Matrix gene (923 nucleotides) | Consistent with A/blue-winged teal/Illinois/10OS1546/2010 (H3N6) Consistent with A/blue-winged teal/Illinois/10OS1546/2010 (H3N6) | 100% identity with A/blue-winged teal/Illinois/ 10OS1546/2010 (H3N6) (GenBank: CY132657) 100% identity with A/blue-winged teal/Illinois/ 10OS1546/2010 (H3N6) (GenBank: CY132658) |
| Titer by CEID₅₀ Assay^{3,4} in Embryonated Chicken Eggs¹ | Report results | 2.0 × 10 ⁸ CEID ₅₀ per mL |
| Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂ | No growth No growth No growth No growth No growth No growth No growth | No growth No growth No growth No growth No growth No growth No growth |
| Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid | None detected None detected | None detected None detected |

¹10-day-old SPF Embryonated Chicken Eggs acquired from B&E Eggs, York Springs, Pennsylvania

²Grown in the allantoic cavity of embryonated chicken eggs¹ for 2 days at 35°C in a humidified chamber

³The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁴2 days at 35°C in a humidified chamber

⁵Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 25 JUL 2013

Signature: 

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

