

***Clostridium cadaveris*, Strain CC88A**

Catalog No. HM-1041

Product Description: *Clostridium cadaveris* (*C. cadaveris*), strain CC88A was isolated in October 2010 from colonic biopsy tissue of a human subject in Victoria, British Columbia, Canada.

Lot^{1,2}: 63266126

Manufacturing Date: 03FEB2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ⁴ Motility (wet mount)	Gram-positive rods Report results Report results	Gram-negative rods ³ Irregular, low convex, undulate, smooth and gray (Figure 1) Motile
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1460 base pairs)	≥ 99% identical to depositor's sequence Consistent with <i>C. cadaveris</i>	≥ 99% identical to depositor's sequence Consistent with <i>C. cadaveris</i>
Purity (post-freeze) Anaerobic growth ⁵ Aerobic growth ⁶	Growth consistent with <i>C. cadaveris</i> No growth	Growth consistent with <i>C. cadaveris</i> No growth
Viability (post-freeze)⁴	Growth	Growth

¹Quality control of HMP material is only performed to demonstrate that the material distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited organism.

²*C. cadaveris*, strain CC88A was deposited by Professor Emma Allen-Vercoe, Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada. The deposited material was inoculated into Modified Chopped Meat media and incubated for 71 hours at 37°C in an anaerobic atmosphere (< 5% O₂; Remel™ Pack-Anaero™). Broth inoculum was added to Tryptic Soy agar with 5% defibrinated sheep blood kolles which were grown 26 hours at 37°C in an anaerobic atmosphere to produce this lot.


³*Clostridium cadaveris* is a Gram-positive organism; however, some species of *Clostridium* which have Gram-positive cell walls will stain Gram-negative or Gram-variable when older cultures (> 24 hours) are used for staining. For additional information, please refer to Beveridge, T. J. "Mechanism of Gram Variability in Select Bacteria." *J. Bacteriol.* 172 (1990): 1609-1620. PubMed: 1689718.

⁴3 days at 37°C in an anaerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood

⁵Purity of this lot was assessed for 8 days on Tryptic Soy agar with 5% defibrinated sheep blood at 37°C in an anaerobic atmosphere.

⁶Purity of this lot was assessed for 8 days on Tryptic Soy agar with 5% defibrinated sheep blood at 37°C in an aerobic atmosphere with 5% CO₂.

Figure 1



Certificate of Analysis for HM-1041

Date: 14 APR 2015

Signature: 

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

