

Certificate of Analysis for NR-30562

Streptococcus agalactiae, Strain BSU188

Catalog No. NR-30562

Product Description: Streptococcus agalactiae (S. agalactiae), strain BSU188 was isolated between in 2003 by Dr. Nadia Brimil from the vagina of a pregnant adult female in Aachen, Germany.

Lot¹: 63527607 Manufacturing Date: 15MAY2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology	Gram-positive cocci	Gram-positive cocci
Colony morphology ²	Report results	Circular, low convex, entire, smooth and cream (Figure 1)
Hemolysis on blood agar ³	Report results	β-hemolytic
Motility (wet mount)	Report results	Non-motile
Biochemical tests:		
Catalase	Negative	Negative
VITEK [®] MS (MALDI-TOF)	Consistent with S. agalactiae	Consistent with S. agalactiae
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1490 base pairs)	Consistent with S. agalactiae	Consistent with S. agalactiae ⁴
Purity (post-freeze) ⁵	Growth consistent with S. agalactiae	Growth consistent with S. agalactiae
Viability (post-freeze) ²	Growth	Growth

¹NR-30562 was produced by inoculation of the deposited material into Tryptic Soy broth and incubated for 1 day at 37°C in an aerobic atmosphere with 5% CO₂. Broth inoculum was added to Tryptic Soy agar kolles which were grown for 1 day under propagation conditions to produce this lot.

Figure 1



Date: 14 JUL 2015

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²1 day on Tryptic Soy agar under propagation conditions

³¹ day on Tryptic Soy agar with 5% defibrinated sheep blood under propagation conditions

⁴Also consistent with Streptococcus difficilis

⁵Purity of this lot was assessed for 7 days at on Tryptic Soy agar under propagation conditions.