

***Trypanosoma brucei* subsp. *rhodesiense*, Strain KETRI 2537**

Catalog No. NR-46436

Product Description: *Trypanosoma brucei* (*T. brucei*) subsp. *rhodesiense*, strain KETRI 2537 was isolated in 1972 from the blood of a patient in Busoga, Uganda. *T. brucei* subsp. *rhodesiense*, strain KETRI 2537 was obtained by Professor C. J. Bacchi from the Kenya Trypanosomiasis Research Institute (KETRI) strain bank at Mugaga, Kenya.

Lot¹: 63803997

Manufacturing Date: 14OCT2015

TEST	SPECIFICATIONS	RESULTS
Genotyping Sequencing of internal transcribed spacer (ITS) 1, 5.8S ribosomal RNA gene, ITS 2 (~ 1290 base pairs)	Consistent with <i>T. brucei</i>	Consistent with <i>T. brucei</i> ²
Functional Activity by PCR Amplification ITS 1, 5.8S ribosomal RNA gene, ITS 2 ³	~ 1300 base pair amplicon	~ 1300 base pair amplicon
Level of Parasitemia (pre-freeze)⁴	≥ 1 × 10 ⁶ parasites/mL	6.6 × 10 ⁶ parasites/mL
Viability (post-freeze)⁵	Growth in inoculated mouse	Growth in inoculated mouse

¹NR-46436 was produced by inoculation of the deposited material into a BALB/c mouse. Infection was allowed to progress for 4 days until the first peak of parasitemia was reached. Infected blood was collected by orbital bleeding and used to inoculate six BALB/c mice. Infection was allowed to progress for 5 days until the first peak of parasitemia was reached and infected blood was collected by orbital bleeding.

²Also consistent with *T. evansi* and/or *T. equiperdum*, which are putative subspecies of *T. brucei* (Lun, Z.-R., et al. "Trypanosoma brucei: Two Steps to Spread Out from Africa." *Trends Parasitol.* 26 (2010): 424-427. PubMed: 20561822.)

³PCR was performed as described in Agbo, E. C., et al. "Measure of Molecular Diversity within the *Trypanosoma brucei* Subspecies *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* as Revealed by Genotypic Characterization." *Exp. Parasitol.* 99 (2001): 123-131. PubMed: 11846522.

⁴Parasitemia was determined after 5 days of infection by microscopic counts using a haemocytometer and 0.85% ammonium chloride as diluent.

⁵Viability of trypanosomes was confirmed by examination of a BALB/c mouse for parasitemia for 7 days.

Date: 26 JAN 2016

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

