

Certificate of Analysis for NR-30649

Mycobacterium tuberculosis, Strain 96-2357

Catalog No. NR-30649

This reagent is the tangible property of the U.S. Government.

Product Description: *Mycobacterium tuberculosis* (*M. tuberculosis*), strain 96-2357 was isolated between 1995 and 2000 from human sputum from an HIV-negative patient infected with pulmonary tuberculosis in North America. Strain 96-2357 was deposited as a multi-drug sensitive (MDS) strain of tuberculosis with sensitivity to rifampicin and isoniazid.

Lot¹: 63103782 Manufacturing Date: 12FEB2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis ²		
Cellular morphology	Gram-positive rods	Gram-positive rods
Colony morphology ³	Report results	Irregular, slight peaked, undulate, rough and cream (Figure 1)
Growth rate	≥ 7 days	19 days
Growth at 26°C	Negative	Negative
Growth at 37°C	Positive	Positive
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment)	Positive (no pigment)	Positive (no pigment)
Biochemical tests		
Niacin production⁴	Positive	Positive
Nitrate reduction	Positive	Positive
Pyrazinamidase	Positive	Positive
Genotypic Analysis		
Sequencing of Heat Shock Protein 65 gene (~ 370 base pairs)	Consistent with M. tuberculosis	Consistent with <i>M. tuberculosis</i> ⁵
Purity (post-freeze) ⁶	Consistent with expected colony morphology	Consistent with expected colony morphology
Viability (post-freeze) ³	Growth	Growth

NR-30649 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment and grown for 35 days at 37°C in an aerobic atmosphere with 5% CO₂. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 20 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org

Fax: 703-365-2898 Tel: 800-359-7370

²Information on *Mycobacterium* testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria." <u>Biochemical Testing</u>. (2012) Jose C. Jimenez-Lopez (Ed.), InTech, http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus *Mycobacterium* and for Description of New Slowly Growing *Mycobacterium* Species." Int. J. Syst. Bacteriol. 42 (1992): 315-323. PubMed: 1581193.

³19 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only *M. tuberculosis* accumulates it, resulting in a positive test for *M. tuberculosis*.

⁵Also consistent with *M. africanum*, *M. bovis*, *M. canettii* and *M. microti*

⁶Purity of this lot was assessed for 21 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment and Tryptic Soy agar plates.



Certificate of Analysis for NR-30649

Figure 1: Colony Morphology



Date: 03 FEB 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®s knowledge.

 ${\sf ATCC}^{\$}$ is a trademark of the American Type Culture Collection. You are authorized to use this product for research use only. It is not intended for human use.