

Certificate of Analysis for NR-41309

Sporothrix globosa, Isolate 8

Catalog No. NR-41309


Product Description: Sporothrix globosa (S. globosa), isolate 8 was obtained from human skin tissue in China in March 2011.

Lot¹: 61697594 Manufacturing Date: 30MAY2013

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Morphology ²	Report results	Ovoid with dark brown pigmentation, undifferentiated hyphae (Figure 1)
Raffinose assimilation ³	Negative	Negative
Genotypic Analysis		
Partial sequencing of internal transcribed spacer (ITS) 1, 5.8S ribosomal RNA (rRNA) gene,	Consistent with S. globosa	Consistent with S. globosa ⁴
and ITS 2 (~ 520 base pairs) Sequencing of beta-tubulin gene	Consistent with S. globosa	Consistent with <i>S. globosa</i> ⁵
(~ 380 base pairs)		
Purity ⁶		
Potato Dextrose agar at 25°C	No bacterial growth	No bacterial growth
Nutrient broth with 0.1% Yeast Extract at 25°C	No bacterial growth	No bacterial growth
Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth	No bacterial growth
Viability (post-freeze) ²	Growth	Growth

NR-41309 was produced by growth of the deposited material on Yeast Mold agar in an aerobic atmosphere at 25°C. An inoculum from the plate was added to Yeast Mold broth and grown 5 days at 25°C in an aerobic atmosphere to produce this lot.

Figure 1: Morphology



Date: 19 FEB 2016 **Signature:**

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²3 days at 25°C in an aerobic atmosphere on Potato Dextrose agar

³5 days at 25°C in an aerobic atmosphere. Nocardia Purple broth with raffinose differentiates *S. globosa* from *S. schenckii* based on the ability of *S. schenckii* isolates to utilize raffinose as a sole carbon source, resulting in a yellow color. *S. globosa* isolates will remain purple. [Marimon, R., et al. "*Sporothrix basiliensis*, *S. globosa*, and *S. mexicana*, Three New *Sporothrix* Species of Clinical Interest." <u>J. Clin. Microbiol.</u> 45 (2007): 3198-3206. PubMed: 17687013.]. Positive control tube (ATCC[®] 20282[™]) was yellow.

⁴99.8% identical to *S. globosa*, type strain CBS 120340 (GenBank: KP017084.1)

⁵99.8% identical to *S. globosa*, strain CBS 129724 (GenBank: KC113238.1), type strain sequence is not available, also consistent with *S. schenckii* ⁶Clarity of broth and lack of growth on agar was determined by visual inspection after 3 days at 25°C and 37°C in an aerobic atmosphere.