

Certificate of Analysis for NR-49439

Influenza A Virus, A/Jiangxi-Donghu/346/2013 (HA, NA) x A/Puerto Rico/8/1934 (H10N8)

Catalog No. NR-49439

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs¹ infected with reassortant influenza A virus, A/Jiangxi-Donghu/346/2013 (HA, NA) x A/Puerto Rico/8/1934 (H10N8)

Lot²: 63777209 Manufacturing Date: 24SEP2015

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs ¹		
Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin and Matrix Coding Regions		
Hemagglutinin (400 nucleotides)	Consistent with A/Jiangxi- Donghu/346/2013 (H10N8)	99% identity with A/Jiangxi- Donghu/346/2013 (H10N8) (EpiFlu: EPI497477) ³
Matrix (919 nucleotides)	Consistent with A/Puerto Rico/8/1934 (H1N1)	100% identity with A/Puerto Rico/8/1934 (H1N1) (GenBank: AF389121)
Titer by CEID ₅₀ Assay ^{4,5} in Embryonated Chicken Eggs ¹	Report results	1.6 × 10 ⁸ CEID ₅₀ per mL
Sterility (21-day incubation)		
Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic	No growth	No growth
Trypticase soy broth, 37°C and 26°C, aerobic	No growth	No growth
Sabouraud broth, 37°C and 26°C, aerobic	No growth	No growth
Blood agar, 37°C, aerobic	No growth	No growth
Blood agar, 37°C, anaerobic	No growth	No growth
Thioglycollate broth, 37°C, anaerobic	No growth	No growth
DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination		
Agar and broth culture (14-day incubation at 37°C)	None detected	None detected
DNA detection by PCR of extracted Test Article nucleic acid	None detected	None detected

¹⁰⁻ to 11-day-old SPF Embryonated Chicken Eggs acquired from B&E Eggs, York Springs, Pennsylvania

Date: 25 MAR 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²Grown in the allantoic cavity of embryonated chicken eggs¹ for 2 days at 35°C in a humidified chamber

³The hemagglutinin gene sequence of influenza A virus, A/Jiangxi-Donghu/346/2013 (H10N8) is identical to GenBank: KJ406543 (influenza A virus, A/Jiangxi-Donghu/IPB13/2013 (H10N8), which was independently isolated from the same patient).

⁴The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁵2 days at 35°C in a humidified chamber

⁶Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.