

Certificate of Analysis for NR-30680

Mycobacterium tuberculosis, Strain 96-2611

Catalog No. NR-30680

This reagent is the tangible property of the U.S. Government.

Product Description: Mycobacterium tuberculosis (M. tuberculosis), strain 96-2611 was isolated between 1995 and 2000 from human sputum from an HIV-negative patient infected with pulmonary tuberculosis in North America. Strain 96-2611 was deposited as a multi-drug sensitive (MDS) strain of tuberculosis with sensitivity to rifampicin and isoniazid.

Lot1: 63385526 Manufacturing Date: 10JUN2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis ²		
Cellular morphology	Gram-positive rods	Gram-positive rods
Colony morphology ³	Report results	Irregular, low convex, undulate, rough and cream (Figure 1)
Growth rate	≥ 7 days	21 days
Growth at 26°C	Negative	Negative
Growth at 37°C	Positive	Positive
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment)	Positive (no pigment)	Positive (no pigment)
Biochemical tests		
Niacin production⁴	Positive	Positive
Nitrate reduction	Positive	Positive
Pyrazinamidase	Positive	Positive
Genotypic Analysis Sequencing of Heat Shock Protein 65 gene (~ 430 base pairs)	≥ 99% sequence identity to <i>M. tuberculosis</i> type strain (GenBank: AL123456)	100% sequence identity to M. tuberculosis type strain (GenBank: AL123456) ⁵
Purity (post-freeze) ⁶	Consistent with expected colony morphology	Consistent with expected colony morphology
Viability (post-freeze) ³	Growth	Growth

NR-30680 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 21 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org Fax: 703-365-2898

Tel: 800-359-7370

²Information on Mycobacterium testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria." <u>Biochemical Testing</u>. (2012) Jose C. Jimenez-Lopez (Ed.), InTech, http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-ofmycobacteria and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus Mycobacterium and for Description of New Slowly Growing *Mycobacterium* Species." Int. J. Syst. Bacteriol. 42 (1992): 315-323. PubMed: 1581193. 321 days at 37°C in an aerobic atmosphere with 5% CO2 on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only *M. tuberculosis* accumulates it, resulting in a positive test for *M. tuberculosis*.

⁵Also consistent with *M. africanum*, *M. bovis*, *M. canettii* and *M. microti*

⁶Purity of this lot was assessed for 36 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment and 21 days at 37°C in an aerobic atmosphere with 5% CO₂ on Tryptic Soy agar plates.



Certificate of Analysis for NR-30680

Figure 1: Colony Morphology



Date: 18 APR 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®s knowledge.

 $ATCC^{\circ}$ is a trademark of the American Type Culture Collection. You are authorized to use this product for research use only. It is not intended for human use.