

N8 Neuraminidase (NA) Protein with N-Terminal Histidine Tag from Influenza Virus, A/chicken/Netherlands/14015531/2014 (H5N8), Recombinant from Baculovirus

Catalog No. NR-50111

This reagent is the tangible property of the U.S. Government.

Product Description: A recombinant form of the N8 neuraminidase (NA) protein from influenza A virus, A/chicken/Netherlands/14015531/2014 (H5N8) containing an N-terminal histidine tag was produced in Sf9 insect cells using a baculovirus expression vector system and was purified by nickel affinity chromatography.

Lot: 64193297

Manufacturing Date: 10MAY2016

TEST	SPECIFICATIONS	RESULTS
Appearance	Clear and colorless	Clear and colorless
SDS-PAGE	Protein band of interest represents > 95% of total staining intensity	Dominant band of ~ 50 kDa accounts for ~ 95% of total staining intensity (Figure 1)
Identification by Western Blot Analysis Polyclonal anti-N8 NA ¹ Monoclonal anti-histidine tag ²	Reactive Reactive	Reactive (Figure 2) Reactive (Figure 3)
Concentration by Bradford Assay³	Report results	152 µg per mL
Final Product Quantity per vial Volume per vial	Report results Report results	79 µg 520 µL
Functional Activity Neuraminidase activity in fluorescent enzymatic assay	Report results	5.3 × 10 ¹⁰ relative fluorescence units per hour per mg protein ⁴
Endotoxin Content (Limulus Amoebocyte Lysate Assay)	Report results	24 EU per mg
Filtration	0.2 µm sterile-filtered	0.2 µm sterile-filtered


¹BEI Resources NR-3145, Polyclonal Anti-Influenza Virus N8 (Neq2) Neuraminidase (NA), A/equine/Miami/1/63 (H3N8), (antiserum, Goat) (1:1000 dilution)

²Clontech 631212 (IgG2a) (1:1000 dilution)

³Using BSA as a standard

⁴Using serial dilutions of NR-50111 and 0.15 mM 2'-(4-methylumbelliferyl)-α-D-N-acetylneuraminic acid (4-MUNANA), Sigma (Cat. No. M8639), as described in Wetherall, N.T., et al. "Evaluation of Neuraminidase Enzyme Assays Using Different Substrates to Measure Susceptibility of Influenza Virus Clinical Isolates to Neuraminidase Inhibitors: Report of the Neuraminidase Inhibitor Susceptibility Network." *J. Clin. Microbiol.* 41 (2003): 742-750. PubMed: 12574276.

Date: 14 JUL 2016

Signature: 

BEI Resources Authentication


ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.




Figure 1: SDS-PAGE




Lane 1: SeeBlue® Plus2 Pre-Stained Protein Standard
 Lane 2: NR-50111, 1.5 µg
 Lane 3: BSA, 2.0 µg

Figure 2: Western Blot with Polyclonal Anti-N8 NA



Lane 1: MagicMark™ XP Protein Standard
 Lane 2: BSA, 0.5 µg
 Lane 3: NR-50111, 0.4 µg

Figure 3: Western Blot with Monoclonal Anti-Histidine Tag



Lane 1: MagicMark™ XP Protein Standard
 Lane 2: BSA, 0.5 µg
 Lane 3: NR-50111, 0.4 µg