

Certificate of Analysis for NR-19017

Mycobacterium tuberculosis, Strain NHN1691

Catalog No. NR-19017

Product Description: *Mycobacterium tuberculosis* (*M. tuberculosis*), strain NHN1691 was isolated in China. Strain NHN1691 was deposited as a multi-drug resistant (MDR) strain of tuberculosis with resistance to rifampicin, isoniazid, ethambutol and streptomycin.

Lot¹: 63383511 Manufacturing Date: 26JUN2015

| TEST | SPECIFICATIONS | RESULTS |
|--|---|---|
| Phenotypic Analysis ² | | |
| Cellular morphology | Gram-positive rods | Gram-positive rods |
| Colony morphology ³ | Report results | Irregular, slight peaked, undulate, rough, opaque and cream (Figure 1) |
| Growth rate | ≥ 7 days | 21 days |
| Growth at 26°C | Negative | Negative |
| Growth at 37°C | Positive | Positive |
| Acid-fast stain | Positive (red colonies) | Positive (red colonies) |
| Pigmentation in the dark (Scotochromogen) | Negative (no pigment) | Negative (no pigment) |
| Photoinduction for 1 hour (Photochromogen) | Negative (no pigment) | Negative (no pigment) |
| Nonchromogen (no pigment) | Positive (no pigment) | Positive (no pigment) |
| Biochemical tests | | |
| Niacin production ⁴ | Positive | Positive |
| Nitrate reduction | Positive | Positive |
| Pyrazinamidase | Positive | Positive |
| Genotypic Analysis Sequencing of Heat Shock Protein 65 gene (~ 380 base pairs) | ≥ 99% sequence identity to M. tuberculosis type strain (GenBank: AL123456) | 100% sequence identity to <i>M. tuberculosis</i> type strain (GenBank: AL123456) ⁵ |
| Purity (post-freeze) ⁶ | Consistent with expected colony morphology | Consistent with expected colony morphology |
| Viability (post-freeze) ³ | Growth | Growth |

¹NR-19017 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles which were grown for 30 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²Information on Mycobacterium testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria" <u>Biochemical Testing</u>. (2012) Jose C. Jimenez-Lopez (Ed.), InTech, http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus *Mycobacterium* and for Description of New Slowly Growing *Mycobacterium* Species." Int. J. Syst. Bacteriol. 42 (1992): 315-323. PubMed: 1581193.

³21 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only *M. tuberculosis* accumulates it, resulting in a positive test for *M. tuberculosis*.

⁵Also consistent with *M. africanum*, *M. bovis*, *M. canettii* and *M. microti*

⁶Purity of this lot was assessed for 66 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment and on Tryptic Soy agar plates.



Certificate of Analysis for NR-19017

Figure 1: Colony Morphology



Date: 11 OCT 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

NR-19017 63383511 11OCT2016